arXiv:1303.2838v1 [math.AP] 12 Mar 2013

Energy decay for a transmission coupled Euler-Bernoulli and wave equation with force and moment feedback

FATHI HASSINE
UR Analysis and Control of PDE (05/UR/15-01)
Département de Mathématiques, Faculté des sciences de Monastir,
5019 Monastir, Tunisie
email: `fathi.hassine@fsm.rnu.tn`

Abstract

In this paper we consider the Euler-Bernoulli equation coupled by the interface with a wave equation and with a boundary damping. The dissipation is acting through the Euler-Bernoulli equation by a force and a moment feedback. We show that in this case, through the Carleman estimates technique that the coupled system is at least logarithmically stable as the time variable goes to infinity. The result follow from the combination of two Carleman estimates: The first is gived for a classical transmission problem while the second one is dedicated to a coupled system where the states are linked on the board.

Key words and phrases: Transmission problem, boundary stabilization, Bernoulli-Euler plate equation, wave equation, logarithmic energy decay, Carleman Estimates.
Mathematics Subject Classification: *35A01, 35A02, 35M32, 35S05, 93D15.*

Contents

1 Introduction and statment of results **2**

2 Existence and uniquness of solution **4**

3 Local Carleman estimate in the surface **7**
  3.1 Reduction of the problem 7
  3.2 Study of the eigenvalues 10
  3.3 Estimation in the region $\mathcal{E}^{+,+}$ 11
  3.4 Estimation in the region $\mathcal{E}^{+,-}$ 16
  3.5 Estimation in the neighborhood of the region $\mathcal{Z}^{+}$ 17
  3.6 Boundary Carleman estimate 21

4 Global Carleman estimates and construction of the weight functions **22**

5 Resolvent Estimate **24**

A Appendix: Semi Classical analysis **30**

References **32**

1 Introduction and statment of results

Let $\Omega = \Omega_1 \cup \Omega_2$, be an open bounded domain in $\mathbb{R}^n$ such that Ω_1 and Ω_2 are two disjoint open connected domains with smooth boundary and verifying that $\overline{\Omega}_1 \cap \overline{\Omega}_2 = S$, $\partial\Omega_1 = S \cup \Gamma_1$ and $\partial\Omega_2 = S \cup \Gamma_2$ where $S \neq \emptyset$, $\Gamma_1 \neq \emptyset$ and $\Gamma_2 \neq \emptyset$ such that $\overline{S} \cap \overline{\Gamma}_1 = \emptyset$ and $\overline{S} \cap \overline{\Gamma}_2 = \emptyset$ (See Figure 1).



Figure 1: Geometrical situation of the transmission Plate/Wave equations.

We Consider the following transmission Plate/Wave system

$$
\begin{cases}
\partial_t^2 u_1 + \Delta^2 u_1 = 0 & \text{in} \quad \Omega_1 \times]0, +\infty[, \\
\partial_t^2 u_2 - \Delta u_2 = 0 & \text{in} \quad \Omega_2 \times]0, +\infty[, \\
u_1 + u_2 = 0 & \text{on} \quad S \times]0, +\infty[, \\
\partial_\nu \Delta u_1 = \partial_\nu u_2 & \text{on} \quad S \times]0, +\infty[, \\
\Delta u_1 = 0 & \text{on} \quad S \times]0, +\infty[, \\
\Delta u_1 = -a \partial_t \partial_\nu u_1 & \text{on} \quad \Gamma_1 \times]0, +\infty[, \\
\partial_\nu \Delta u_1 = b \partial_t u_1 & \text{on} \quad \Gamma_1 \times]0, +\infty[, \\
u_2 = 0 & \text{on} \quad \Gamma_2 \times]0, +\infty[, \\
u_1(x,0) = u_1^0(x),\ \partial_t u_1(x,0) = u_1^1(x) & \text{in} \quad \Omega_1, \\
u_2(x,0) = u_2^0(x),\ \partial_t u_2(x,0) = u_2^1(x) & \text{in} \quad \Omega_2.
\end{cases} \tag{1.1}
$$

Where ν denotes the outward normal vector of Ω_1 in Γ_1 and S and of Ω_2 in Γ_2 only; and a and b are two non negative bounded functions on Γ_1 such that there exists a strictly positive constant c_0 verifying

$$
a \geq c_0 \quad \text{and} \quad b \geq c_0 \quad \text{on } \Gamma_1. \tag{1.2}
$$

We define the energy of a solution

$$
u = \begin{cases} u_1 & \text{in} \quad \Omega_1 \\ u_2 & \text{in} \quad \Omega_2 \end{cases}
$$

of the system (1.1) at the time $t \geq 0$ by

$$
E(t,u) = \frac{1}{2}\left(\int_{\Omega_1} |\partial_t u_1(x,t)|^2 + |\Delta u_1(x,t)|^2 \,\mathrm{d}x + \int_{\Omega_2} |\partial_t u_2(x,t)|^2 + |\nabla u_2(x,t)|^2 \,\mathrm{d}x \right).
$$

By means of the classical energy method, we have

$$
E(t_2,u) - E(t_1,u) = -\int_{t_1}^{t_2}\!\!\int_{\Gamma_1} a |\partial_\nu \partial_t u_1(x,t)|^2 \,\mathrm{d}x\,\mathrm{d}t - \int_{t_1}^{t_2}\!\!\int_{\Gamma_1} b |\partial_t u_1(x,t)|^2 \,\mathrm{d}x\,\mathrm{d}t, \quad \forall\, t_1, t_2 > 0,
$$

this means that the energy dissipates over time and in this paper we are going to measure this dissipation rate.

We define the Hilbert space $\mathcal{H} = X \times H$ where $H = L^2(\Omega_1) \times L^2(\Omega_2)$ and

$$X = \Big\{u = (u_1, u_2) \in Y;\ u_1 \in H^2(\Omega_1),\ u_{1\,|S} + u_{2\,|S} = 0,\ u_{2\,|\Gamma_2} = 0,\\ \int_S u_1.\overline{\partial_\nu u_1}\mathrm{d}x + \int_{\Gamma_1} u_1.\overline{\partial_\nu u_1}\mathrm{d}x = 0\Big\},$$

with $Y = L^2(\Omega_1) \times H^1(\Omega_2)$. The space $\mathcal{H}$ is equipped with the norm

$$\|(u,v)\|^2_{\mathcal{H}} = \|\Delta u_1\|^2_{L^2(\Omega_1)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} + \|v_1\|^2_{L^2(\Omega_1)} + \|v_2\|^2_{L^2(\Omega_2)},$$

where we denote by

$$u = \left\{\begin{array}{lll} u_1 & \text{in} & \Omega_1 \\ u_2 & \text{in} & \Omega_2 \end{array}\right. \quad \text{and} \quad v = \left\{\begin{array}{lll} v_1 & \text{in} & \Omega_1 \\ v_2 & \text{in} & \Omega_2. \end{array}\right.$$

We define the operator

$$A = \begin{bmatrix} 0 & 0 & \mathrm{Id} & 0 \\ 0 & 0 & 0 & \mathrm{Id} \\ -\Delta^2 & 0 & 0 & 0 \\ 0 & \Delta & 0 & 0 \end{bmatrix}$$

whose domain is gived by

$$\mathcal{D}(\mathcal{A}) = \big\{(u,v) = (u_1,u_2,v_1,v_2) \in \mathcal{H} : (v_1,v_2,\Delta^2 u_1, \Delta u_2) \in \mathcal{H},\ \partial_\nu \Delta u_{1\,|S} = \partial_\nu u_{2\,|S},\\ \Delta u_{1\,|S} = 0, \Delta u_{1\,|\Gamma_1} = -a\partial_\nu v_{1\,|\Gamma_1},\ \partial_\nu \Delta u_{1\,|\Gamma_1} = b v_{1\,|\Gamma_1}\big\}.$$

Our main result is the following

Theorem 1.1 *For any $k \in \mathbb{N}$ there exists a constant $C > 0$ such that for any initial data $(u^0, u^1) = (u^0_0, u^0_1, u^1_0, u^1_1) \in \mathcal{D}(\mathcal{A}^k)$ the solution $u(x,t)$ of (1.1) starting from (u^0, u^1) satisfying*

$$E(t,u) \leq \frac{C}{(\ln(2+t))^{2k}} \|(u^0, u^1)\|^2_{\mathcal{D}(\mathcal{A}^k)}.$$

This theorem is resulting, by Burq's [Bur98], from the resolvent estimate gived by the following theorem

Theorem 1.2 *There exists $C > 0$ such that for every $\mu \in \mathbb{R}$ with $|\mu|$ large, we have*

$$\|(\mathcal{A} - i\mu\,\mathrm{Id})^{-1}\|_{\mathcal{L}(\mathcal{H})} \leq C\mathrm{e}^{C|\mu|}.$$

Several authors are intersted to the decay rate of energy for transmission problems:

- For transmission wave equation with diffusion coefficient, Bellassoued [Bel03] proved that the local energy decreasing logarithmically by the means of Carleman estimates.
- For a hyperbolic-parbolic coupled systems arising in fluide structure interaction, It's proved, under geometric control conditions, by Rauch, Zhang and Zuazua [RZZ05], Zhang and Zuazua in [ZZ06] and [ZZ07] and by Duyckaerts [Duy07] that we have a polynomial decay of a solution and they proved also the lack of uniform decay of enregy. Without the geometric control condition it was proved that the energy is deceasing logarithmically by Zhang and Zuazua [ZZ06] by means of logarithmic observability estimate, and by Duyckaerts [Duy07] by the means of Carleman estimates both for non-natural transmission condition, and by Fathallah [Fat11] for natural transmission condition by the means of Carleman estimates.
- For the Bernoulli-Euler plate equation Ammari and Vodev [AV09] proved that the uniform decay of solution, and for different geometric domain with force and moment Hassine [Has13] proved a logarithmic decay of energy by the means of Carleman estimates.

Tibou [Teb12] has interested to the coupled Euler-Bernoulli and wave equations without transmission and he proved the polynomial decay of energy by the means of multiplier method, for two models whose the first is where the dissipation acts through the Euler-Bernoulli equation and the second one is acting through the wave equation. For a transmission model, Ammari and Nicaise [AN10] have proved, under some geometric condition, an exponential stabilization for a coupled damped wave equation with a damped Kirchhoff plate equation.

In this paper we will prove that a solution of the system (1.1) is decreasing at least logarithmically along the time. Inspired from [Has13], to prove the resolvent estimate gived in Theorem 1.2, we will combine two suitable Carleman estimates whose the first is applyed to a system of second ordre with transmission conditions and the second one is applyed to a system of second order also in Ω_1 with two operators and two states linked by boundary conditions at the interface S.

The outline of this paper is as follows: In section 2 we prove the well-posedness of the problem (1.1), in section 3 we will introduce the local Carleman estimates, in section 4 we give the global Carleman estimates and we construct a suitable weight functions that satisfying the Hörmander's assumption, in section 5 we prove the resolvent estimate and finally in Appendix A we will give same notations and some important results of microlocal analysis used essentially in section 3.

In these notes C will always be a generic positive constant whose value may be different from one line to another, and where the constant depends on a number of variables is denoted indexed by them.

2 Existence and uniquness of solution

In this section we will prove the Well-posedness of the problem (1.1) through the semi-classical analysis. The system (1.1) can be writen in the abstract form as a Cauchy problem as follows

$$\begin{cases} \partial_t \begin{pmatrix} u_1 \\ u_2 \\ v_1 \\ v_2 \end{pmatrix}(t,x) = \mathcal{A}\begin{pmatrix} u_1 \\ u_2 \\ v_1 \\ v_2 \end{pmatrix}(t,x) & \text{if} \quad (t,x) \in]0,+\infty[\times\Omega, \\ \begin{pmatrix} u_1 \\ u_2 \\ v_1 \\ v_2 \end{pmatrix}(0,x) = \begin{pmatrix} u_1^0 \\ u_2^0 \\ u_1^1 \\ u_2^1 \end{pmatrix}(x) & \text{if} \quad x \in \Omega, \end{cases}$$

where we recall that the operator $\mathcal{A}$ is defined by

$$A = \begin{bmatrix} 0 & 0 & \mathrm{Id} & 0 \\ 0 & 0 & 0 & \mathrm{Id} \\ -\Delta^2 & 0 & 0 & 0 \\ 0 & \Delta & 0 & 0 \end{bmatrix}$$

with domain

$$\begin{aligned} \mathcal{D}(\mathcal{A}) = \big\{(u,v) = (u_1,u_2,v_1,v_2) \in \mathcal{H} : (v_1,v_2,\Delta^2 u_1, \Delta u_2) \in \mathcal{H},\ \partial_\nu \Delta u_{1\,|S} = \partial_\nu u_{2\,|S}, \\ \Delta u_{1\,|S} = 0, \Delta u_{1\,|\Gamma_1} = -a\partial_\nu v_{1\,|\Gamma_1},\ \partial_\nu \Delta u_{1\,|\Gamma_1} = b v_{1\,|\Gamma_1}\big\}. \end{aligned}$$

In the space $Y = L^2(\Omega_1) \times H^1(\Omega_2)$ we define the operator G as follows

$$Gu = G\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} -\Delta u_1 \\ u_2 \end{pmatrix}$$

with domain

$$\begin{aligned} D(G) = X = \Big\{u = (u_1,u_2) \in Y;\ u_1 \in H^2(\Omega_1),\ u_{1\,|S} + u_{2\,|S} = 0,\ u_{2\,|\Gamma_2} = 0, \\ \int_S u_1.\overline{\partial_\nu u_1}\mathrm{d}x + \int_{\Gamma_1} u_1.\overline{\partial_\nu u_1}\mathrm{d}x = 0\Big\}, \end{aligned}$$

We define a norm in the space X by

$$\|u\|_X^2 = \|(u_1, u_2)\|_X^2 = \|\Delta u_1\|^2_{L^2(\Omega_1)} + \|\nabla u_2\|^2_{L^2(\Omega_2)},$$

and we recall the graph norm of G is gived by

$$\|u\|^2_{gr(G)} = \|u_1\|^2_{L^2(\Omega_1)} + \|\Delta u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L^2(\Omega_2)} + \|\nabla u_2\|^2_{L^2(\Omega_2)}.$$

Lemma 2.1 $(X, \| \,.\, \|_X)$ *is a Hilbert space with a norm equivalent to the graph norm of* G*.*

Proof :
It is easy to show that by Green's formula and Poincaré inequality to show that there exists $C > 0$ such that for every $u = (u_1, u_2) \in X$ we have

$$\langle Gu, u\rangle_Y = \|\nabla u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L^2(\Omega_2)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} \geq C(\|u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L^2(\Omega_2)} + \|\nabla u_2\|^2_{L^2(\Omega_2)}).$$

Then G is a strictly positive operator in Y and for every $u = (u_1, u_2) \in X$ we have

$$\|u\|_X.\|u\|_Y \geq C\|Gu\|_Y.\|u\|_Y \geq C\langle Gu, u\rangle_H \geq C\|u\|^2_Y,$$

which prove the equivalence the two norms.
Since G is positive then by [TW09, Proposition 3.3.5] $-G$ is m-dissipatif in particular G is a closed operator, and this conclude the proof while $(X, \| \,.\, \|_{gr(G)})$ is now a Hilbert space. ■

Remarks 2.1 *1. The two norms* $\| \,.\, \|_{H^2(\Omega_1)} + \|\nabla . \|_{L^2(\Omega_2)}$ *and* $\| \,.\, \|_X$ *are equivalent in* X*, in fact* $(X, \| \,.\, \|_{H^2(\Omega_1)} + \|\nabla . \|_{L^2(\Omega_2)})$ *is a Hilbert space while* X *is a closed subset of* $H^2(\Omega_1) \times H^1(\Omega_2)$ *and we have*

$$\|u\|_X \leq C(\|u_1\|_{H^2(\Omega_1)} + \|u_2\|_{H^1(\Omega_2)}), \quad \forall\, u = (u_1, u_2) \in X,$$

then by Banach theorem [EMT04, p.132] we obtain easly the result.

2. By combination of Lemma 2.1 and the previous remark we follow that X *is also a Hilbert space with the norm*

$$\| \,.\, \|_{gr(G)} + \|a^{\frac{1}{2}}\partial_\nu \,.\, \|_{L^2(\Gamma_1)} + \|b^{\frac{1}{2}} \,.\, \|_{L^2(\Gamma_1)}.$$

Theorem 2.1 *The operator* $\mathcal{A}$ *is m-dissipative and especially it generates a strongly semigroup of contraction in* $\mathcal{H}$*.*

Proof :
According to Lumer-Phillips theorem (see for exemple [TW09, p.103]) we have only to prove that $\mathcal{A}$ is m-dissipative.
Let $(u, v) \in \mathcal{D}(\mathcal{A})$ then by Green's formula we have

$$\mathrm{Re}\left\langle \mathcal{A}\begin{pmatrix} u \\ v \end{pmatrix}, \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle_{\mathcal{H}} = \mathrm{Re}\Big(\langle \Delta v_1, \Delta u_1\rangle_{L^2(\Omega_1)} - \langle \Delta^2 u_1, v_1\rangle_{L^2(\Omega_1)} + \langle \Delta u_2, v_2\rangle_{L^2(\Omega_2)}$$
$$+ \langle \nabla v_2, \nabla u_2\rangle_{L^2(\Omega_2)}\Big) = -(\|a^{\frac{1}{2}}\partial_\nu v_1\|^2_{L^2(\Gamma_1)} + \|b^{\frac{1}{2}} v_1\|^2_{L^2(\Gamma_1)}) \leq 0.$$

This shows that $\mathcal{A}$ is dissipative.
Let now $(f, g) \in \mathcal{H}$ and let's find a quadruplet $(u_1, u_2, v_1, v_2) \in \mathcal{D}(\mathcal{A})$ such that

$$(\mathrm{Id} - \mathcal{A})\begin{pmatrix} u_1 \\ u_2 \\ v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} u_1 - v_1 \\ u_2 - v_2 \\ v_1 + \Delta^2 u_1 \\ v_2 - \Delta u_2 \end{pmatrix} = \begin{pmatrix} f_1 \\ f_2 \\ g_1 \\ g_2 \end{pmatrix}$$

then we find that (u_1, u_2, v_1, v_2) satisfy to the following system

$$\left\{\begin{array}{lll} v_1 = u_1 - f_1 & \text{in} & \Omega_1 \\ v_2 = u_2 - f_2 & \text{in} & \Omega_2 \\ u_1 + \Delta^2 u_1 = f_1 + g_1 & \text{in} & \Omega_1 \\ u_2 - \Delta u_2 = f_2 + g_2 & \text{in} & \Omega_2. \end{array}\right.$$

From the second point of Remarks 2.1 and the Riesz representation theorem, there exists a unique $u \in X = \mathcal{D}(G)$ such that for all $\varphi = (\varphi_1, \varphi_2) \in X$ we have

(2.1)
$$\begin{gathered}\langle f_1 + g_1, \varphi_1\rangle_{L^2(\Omega_1)} + \langle f_2 + g_2, \varphi_2\rangle_{L^2(\Omega_2)} + \langle a\partial_\nu f_1, \partial_\nu \varphi_2\rangle_{L^2(\Gamma_1)} + \langle b f_1, \varphi_1\rangle_{L^2(\Gamma_1)} = \langle u_1, \varphi_1\rangle_{L^2(\Omega_1)} + \\ \langle \Delta u_1, \Delta\varphi_1\rangle_{L^2(\Omega_1)} + \langle u_2, \varphi_2\rangle_{L^2(\Omega_2)} + \langle \nabla u_2, \nabla\varphi_2\rangle_{L^2(\Omega_2)} + \langle a\partial_\nu u_1, \partial_\nu\varphi_1\rangle_{L^2(\Gamma_1)} + \langle b u_1, \varphi_1\rangle_{L^2(\Gamma_1)}.\end{gathered}$$

Then by Green's formula we obtain

$$\begin{gathered}\langle f_1 + g_1 - \Delta^2 u_1 - u_1, \varphi_1\rangle_{L^2(\Omega_1)} + \langle f_2 + g_2 + \Delta u_2 - u_2, \varphi_2\rangle_{L^2(\Omega_2)} = \langle \Delta u_1, \partial_\nu\varphi_1\rangle_{L^2(S)} + \\ \langle \partial_\nu u_2 - \partial_\nu\Delta u_1, \varphi_1\rangle_{L^2(S)} + \langle \Delta u_1 + a(\partial_\nu u_1 - \partial_\nu f_1), \partial_\nu\varphi_1\rangle_{L^2(\Gamma_1)} + \langle b(u_1 - f_1) - \partial_\nu\Delta u_1, \varphi_1\rangle_{L^2(\Gamma_1)}.\end{gathered}$$

In particular for $(\varphi_1, \varphi_2) \in \mathscr{C}_c^\infty(\Omega_1) \times \mathscr{C}_c^\infty(\Omega_2)$ we have

$$\langle f_1 + g_1 - \Delta^2 u_1 - u_1, \varphi_1\rangle_{L^2(\Omega_1)} + \langle f_2 + g_2 + \Delta u_2 - u_2, \varphi_2\rangle_{L^2(\Omega_2)} = 0$$

then we find

$$\left\{\begin{array}{ll} u_1 + \Delta^2 u_1 = f_1 + g_1 & \text{in } L^2(\Omega_1) \\ u_2 - \Delta u_2 = f_2 + g_2 & \text{in } L^2(\Omega_2). \end{array}\right. \tag{2.2}$$

Then by (2.1) and (2.2) and (2.2) we obtain

$$\begin{aligned}\langle \Delta u_1, \partial_\nu\varphi_1\rangle_{L^2(S)} + \langle \partial_\nu u_2 - \partial_\nu\Delta u_1, \varphi_1\rangle_{L^2(S)} + \langle \Delta u_1 + a(\partial_\nu u_1 - \partial_\nu f_1), \partial_\nu\varphi_1\rangle_{L^2(\Gamma_1)} \\ + \langle b(u_1 - f_1) - \partial_\nu\Delta u_1, \varphi_1\rangle_{L^2(\Gamma_1)} = 0\end{aligned}$$

and this show the following equalities

$$\Delta u_{1\,|S} = 0,\ \partial_\nu\Delta u_{1\,|S} = \partial_\nu u_{2\,|S},$$

and also

$$\begin{aligned}\Delta u_{1\,|\Gamma_1} &= -a(\partial_\nu u_{1\,|\Gamma_1} - \partial_\nu f_{1\,|\Gamma_1}) = -a\,\partial_\nu v_{1\,|\Gamma_1} \\ \partial_\nu\Delta u_{1\,|\Gamma_1} &= b(u_{1\,|\Gamma_1} - f_{1\,|\Gamma_1}) = b\,v_{1\,|\Gamma_1}.\end{aligned}$$

And this give end to our proof. ∎

One consequence of this last result is that if we assume that $(u_0, v_0) \in \mathcal{D}(\mathcal{A})$, there exists a unique solution of (1.1) which can be expressed by means of a semigroup on $\mathcal{H}$ as follows

$$\begin{pmatrix} u \\ \partial_t u \end{pmatrix} = e^{t\mathcal{A}} \begin{pmatrix} u_0 \\ v_0 \end{pmatrix} \tag{2.3}$$

where $e^{t\mathcal{A}}$ is the semigroupe of the operator $\mathcal{A}$. And we have the following regularity of the solution

$$\begin{pmatrix} u \\ \partial_t u \end{pmatrix} \in C([0, +\infty[, \mathcal{D}(\mathcal{A})) \cap C^1([0, +\infty[, \mathcal{H}).$$

And if $(u_0, v_0) \in \mathcal{H}$, the function $u(t)$ given by (2.3) is the mild solution of (1.1).

3 Local Carleman estimate in the surface

Let $u = (u_1, u_2)$ and $f = (f_1, f_2)$ satisfying the following equations

$$\left\{\begin{array}{ll} A_1(x,D)(u_1) = f_1 & \text{in } \Omega_1 \\ A_2(x,D)(u_2) = f_2 & \text{in } \Omega_2 \\ u_1 = u_2 + \theta & \text{on } S \\ \partial_\nu u_1 = \partial_\nu u_2 + \tilde{\theta} & \text{on } S \end{array}\right. \tag{3.1}$$

where $A_1(x,D)$ and $A_2(x,D)$ defined one differential operator with C^∞-coefficients gived by

$$A(x,D) = \left\{\begin{array}{ll} A_1(x,D)(u_1) = -\Delta u_1 & \text{in } \Omega_1 \\ \\ A_2(x,D)(u_2) = -\Delta u_2 - \dfrac{1}{h^2} u_2 & \text{in } \Omega_2 \end{array}\right.$$

with h is a small pseudo-differential parameter. This mean that the continuity of u at the interface, as well as the continuity of the normal flux, is modulo some error terms θ and $\tilde{\theta}$ respectively. The aim of this section is to find a local Carleman estimate near the interface for a such problem.

3.1 Reduction of the problem

In a neighborhood V of a point $z = (z', z_n) \in S$, we denote by x_n the variable that is normal to the interface and by x' the remainding variables, i.e $x = (x', x_n)$ in particular $z = (z', 0)$. The interface is given now by $S = \{x_n = 0\}$. In a sufficient small neighborhood $V \subset \mathbb{R}^n$ of z we place our selves in normal geodesic coordinates. For convience, we shall take the neighborhood V of the form $V = V_{z'} \times]-\epsilon, \epsilon[$ where $V_{z'}$ is a sufficient small neighborhood of z'. In such coordinate system the operator A take the following form [Hör85, Appendix C] on both side of the interface

$$A(x,D) := \left\{\begin{array}{ll} A_1(x,D) := -\partial^2_{x_n} + R_1(x, \partial_{x'}/i) & x_n > 0 \\ \\ A_2(x,D) := -\partial^2_{x_n} + R_2(x, \partial_{x'}/i) - \dfrac{1}{h^2} & x_n < 0 \end{array}\right.$$

where

$$R(x,\xi') = \left\{\begin{array}{ll} R_1(x,\xi') & x_n > 0 \\ R_2(x,\xi') & x_n < 0 \end{array}\right.$$

is a second ordre polynomial in ξ' with coefficients in $\mathbb{R}$ with principal symbol

$$r(x,\xi') = \left\{\begin{array}{ll} r_1(x,\xi') & x_n > 0 \\ r_2(x,\xi') & x_n < 0 \end{array}\right.$$

that satisfy

$$\left\{\begin{array}{ll} r_1(x,\xi') \geq C|\xi'|^2 & \forall\, (x', x_n) \in V_{z'} \times]0, \epsilon[,\ \forall\, \xi' \in \mathbb{R}^{n-1} \\ r_2(x,\xi') \geq C|\xi'|^2 & \forall\, (x', x_n) \in V_{z'} \times]-\epsilon, 0[,\ \forall\, \xi' \in \mathbb{R}^{n-1}. \end{array}\right. \tag{3.2}$$

Follow to [RR10] we set

$$\mathbb{R}^n_- = \{x :\ x_n < 0\}, \quad \mathbb{R}^n_+ = \{x :\ x_n > 0\}, \quad V^- = V \cap \mathbb{R}^n_-, \quad V^+ = V \cap \mathbb{R}^n_+.$$

For a compact set K of V we set $K^- = K \cap \overline{\mathbb{R}^n_-}$ and $K^+ = K \cap \overline{\mathbb{R}^n_+}$. We then denote by $\mathscr{C}^\infty_c(K^+)$ (resp. $\mathscr{C}^\infty_c(K^-)$) the space of functions that are $\mathscr{C}^\infty$ in $\overline{\mathbb{R}^n_+}$ (resp. $\overline{\mathbb{R}^n_-}$) with support in K^+ (resp. K^-).

We let $\varphi = (\varphi_1, \varphi_2)$ a weight function and we define in both side of S the adjoint operator

$$A_\varphi = h^2 \mathrm{e}^{\varphi/h} A \mathrm{e}^{-\varphi/h}$$

with principal symbol

$$a_\varphi(x,\xi) = \begin{cases} (\xi_n + i(\partial_{x_n}\varphi_1))^2 + r_1(x,\xi' + i(\partial_{x'}\varphi_1)) & x_n > 0 \\ (\xi_n + i(\partial_{x_n}\varphi_2))^2 + r_2(x,\xi' + i(\partial_{x'}\varphi_2)) - 1 & x_n < 0. \end{cases}$$

Assumption 3.1 *The weight function* φ *is in* $\mathscr{C}(\overline{V})$ *and* $\varphi_{|\overline{\mathbb{R}^n_\pm}} \in \mathscr{C}^\infty(\overline{V}^\pm)$ *and satisfies*

1. $|\nabla\varphi|(x) > 0$ *in* $\overline{V}$.
2. $\begin{cases} (\partial_{x_n}\varphi_1)(x',0) > 0 \text{ and } (\partial_{x_n}\varphi_2)(x',0) > 0^1 & \\ (\partial_{x_n}\varphi_1)(x',0) - (\partial_{x_n}\varphi_2)(x',0) > 0 & \text{for all } x' \in V_{z'}. \\ (\partial_{x_n}\varphi_1)^2(x',0) - (\partial_{x_n}\varphi_2)^2(x',0) > 1 & \end{cases}$
3. *The sub-ellipticity condition:*

$$\forall\,(x,\xi) \in \overline{V} \times \mathbb{R}^n;\ a_\varphi(x,\xi) = 0 \Longrightarrow \{\mathrm{Re}(a_\varphi), \mathrm{Im}(a_\varphi)\}(x,\xi) > 0. \tag{3.3}$$

We set

$$f = \begin{cases} f_1 & \text{in } V \cap \Omega_1 \\ f_2 & \text{in } V \cap \Omega_2, \end{cases}$$

then from the assumptions made on the weight function φ we shall obtain the following local Carleman estimate.

Theorem 3.1 *Let* K *be a compact subset of* V *and* φ *a weight function satisfying Assumption 3.1, then there exist* $C > 0$ *and* $h_0 > 0$ *such that*

$$\begin{aligned} &h\|e^{\varphi/h}u\|_0^2 + h^3\|e^{\varphi/h}\nabla u\|_0^2 + h|e^{\varphi/h}u_{|x_n=0}|_0^2 + h^3|e^{\varphi/h}\nabla_{x'}u_{|x_n=0}|_0^2 + h^3|e^{\varphi/h}\partial_{x_n}u_{|x_n=0}|_0^2 \\ &\qquad \le C(h^4\|e^{\varphi/h}f\|_0^2 + h|e^{\varphi/h}\theta|_0^2 + h^3|e^{\varphi/h}\nabla_{x'}\theta|_0^2 + h^3|e^{\varphi/h}\tilde{\theta}|_0^2) \end{aligned}$$

for all $0 < h \le h_0$, u *and* f *satisfying* (3.1) *where* $u_{|\overline{\mathbb{R}^n_\pm}} \in \mathscr{C}^\infty_c(K^\pm)$.

Proof :
We choose χ^+, χ^- and χ^0 with values in $[0,1]$ that satisfy the properties listed in Proposition 3.2, Proposition 3.3 and Proposition 3.4 respectively such that $\chi^+ + \chi^- + \chi^0 \equiv 1$ in a neighborhood of $K \times \mathbb{R}^n$. For $v = e^{\Phi/h}u$ and since $\chi^+ + \chi^- + \chi^0 \equiv 1$ in the neighborhood of $\mathrm{supp}(v)$ we have

$$\begin{cases} |(D_{x_n}v)_{|x_n=0}|_0 \le |\mathrm{op}_t(\chi^+)(D_{x_n}v)_{|x_n=0}|_0 + |\mathrm{op}_t(\chi^-)(D_{x_n}v)_{|x_n=0}|_0 + |\mathrm{op}_t(\chi^0)(D_{x_n}v)_{|x_n=0}|_0 \\ \\ |v_{|x_n=0}|_1 \le |\mathrm{op}_t(\chi^+)v_{|x_n=0}|_1 + |\mathrm{op}_t(\chi^-)v_{|x_n=0}|_1 + |\mathrm{op}_t(\chi^0)v_{|x_n=0}|_1 + Ch|v_{|x_n=0}|_1 \\ \\ \|v\|_1 \le \|\mathrm{op}_t(\chi^+)v\|_1 + \|\mathrm{op}_t(\chi^-)v\|_1 + \|\mathrm{op}_t(\chi^0)v\|_1 + Ch\|v\|_1. \end{cases}$$

Thus if we apply Proposition 3.2, Proposition 3.3 and Proposition 3.4 respectively to the functions χ^+v, χ^-v and χ^0v we obtain for h sufficiently small

$$h\|v\|_1^2 + h|v_{|x_n=0}|_1^2 + h|(D_{x_n}v)_{|x_n=0}|_0^2 \le C(\|P_\varphi v\|_0^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2). \tag{3.4}$$

Observing now that

$$\begin{cases} \|e^{\Phi/h}D_{x_j}u\|_0 \le \|D_{x_j}v\|_0 + \|(\partial_{x_j}\Phi)e^{\Phi/h}u\|_0 \\ |e^{\Phi/h}(D_{x_j}u)_{|x_n=0}|_0 \le |(D_{x_j}v)_{|x_n=0}|_0 + |(\partial_{x_j}\Phi)e^{\Phi/h}u_{|x_n=0}|_0 \end{cases}$$

then the result can be achieved by (3.4) with recalling the forms of θ_φ and $\tilde{\theta}_\varphi$ in (3.8). ■

---

[1] This condition is not necessary as we will see in the sequel but it will help us to describe the position of the roots of some polynomials, ones relative to each others

Follow to [Bel03] we shall consider the transmission problem as a system of two equations in V^+ coupled at the boundary $x_n = 0$. Thus we make the change of variables x_n to $-x_n$ in V^-. This yield the following boundary value problem in V^+:

$$\tag{3.5} \begin{cases} (-\partial^2_{x_n} + R^+(x, \partial_{x'}/i))u^+ = f^+ & x_n > 0 \\ (-\partial^2_{x_n} + R^-(x, \partial_{x'}/i))u^- - \dfrac{1}{h^2}u^- = f^- & x_n > 0 \\ u^+ = u^- + \theta & x_n = 0 \\ \partial_{x_n} u^+ + \partial_{x_n} u^- = \tilde{\theta} & x_n = 0, \end{cases}$$

where for ψ defined in V, we set $\psi^+ = \psi_{|V^+}$ and $\psi^- = \psi_{|V^-}(x', -x_n)$ for $x_n > 0$ and $R^-(x, \partial_{x'}/i) = R(x', -x_n, \partial_{x'}/i)$ and $R^+(x, \partial_{x'}/i) = R(x, \partial_{x'}/i)$ for $x_n > 0$. From that the second point in Assumption 3.1 will be

$$\tag{3.6} \begin{cases} (\partial_{x_n}\varphi^+)_{|x_n=0} > 0 \ ; \ (\partial_{x_n}\varphi^-)_{|x_n=0} < 0 \\ (\partial_{x_n}\varphi^+)_{|x_n=0} + (\partial_{x_n}\varphi^-)_{|x_n=0} > 0 \\ (\partial_{x_n}\varphi^+)^2_{|x_n=0} - (\partial_{x_n}\varphi^-)^2_{|x_n=0} > 1. \end{cases}$$

However the fourth point is preserved since $\{\mathrm{Re}(a_\varphi), \mathrm{Im}(a_\varphi)\}$ is invariant under a change of variables (see [Hör63, p.186]). We set the operator

$$P = \begin{pmatrix} -\partial^2_{x_n} + R^+(x, \partial_{x'}/i) & 0 \\ 0 & -\partial^2_{x_n} + R^-(x, \partial_{x'}/i) - h^{-2} \end{pmatrix}$$

with pricipal symbol

$$p = \begin{pmatrix} \xi_n^2 + r^+(x, \xi') & 0 \\ 0 & \xi_n^2 + r^-(x, \xi') - h^{-2} \end{pmatrix}.$$

For $\Phi = \begin{pmatrix} \varphi^+ & 0 \\ 0 & \varphi^- \end{pmatrix}$, we set $v = {}^t(v^+, v^-)$ for $v = \mathrm{e}^{\Phi/h}u$, then v satisfy the following boundary conditions:

$$\tag{3.7} \begin{cases} v^+_{|x_n=0} = v^-_{|x_n=0} + \theta_\varphi \\ (D_{x_n} + i(\partial_{x_n}\varphi^+))v^+_{|x_n=0} + (D_{x_n} + i(\partial_{x_n}\varphi^-))v^-_{|x_n=0} = \tilde{\theta}_\varphi \end{cases}$$

where

$$\tag{3.8} \theta_\varphi = \theta(\mathrm{e}^{\varphi/h})_{|x_n=0} \quad \text{and} \quad \tilde{\theta}_\varphi = \frac{h}{i}\tilde{\theta}(\mathrm{e}^{\varphi/h})_{|x_n=0}.$$

We define the conjugate operator

$$P_\varphi = h^2 \mathrm{e}^{\Phi/h} P \mathrm{e}^{-\Phi/h}$$

with principal symbol p_φ written in both side as follows

$$p_\varphi(x, \xi) = \begin{cases} p^+_\varphi(x, \xi) = (\xi_n + i\partial_{x_n}(\varphi^+))^2 + r^+(x, \xi' + i\partial_{x'}(\varphi^+)) \\ p^-_\varphi(x, \xi) = (\xi_n + i\partial_{x_n}(\varphi^-))^2 + r^-(x, \xi' + i\partial_{x'}(\varphi^-)) - 1. \end{cases}$$

Separating the real and imaginary part of p_φ, we can write following to [LR95] that

$$\begin{cases} p^+_\varphi = \tilde{q}_2^+ + i\tilde{q}_1^+ \\ p^-_\varphi = \tilde{q}_2^- + i\tilde{q}_1^- \end{cases} \text{where } \begin{cases} \tilde{q}_2^\pm = \xi_n^2 + q_2^\pm \\ \tilde{q}_1^\pm = 2(\partial_{x_n}\varphi^\pm)\xi_n + q_1^\pm \end{cases}$$

with

$$\tag{3.9} \begin{cases} q_2^\pm(x, \xi') = -(\partial_{x_n}\varphi^\pm)^2 + r^\pm(x, \xi') - r^\pm(x, \partial_{x'}\varphi^\pm) & \in S^2_t \\ q_1^\pm(x, \xi') = \tilde{r}^\pm(x, \xi', \partial_{x'}\varphi^\pm) & \in S^1_t, \end{cases}$$

where $\tilde{r}^\pm(x, \xi', \eta') \in S^1_t$ are the symetric bilinear forms in (ξ', η') associated to the real quadratic forms $r^\pm(x, \xi')$ in ξ'.

3.2 Study of the eigenvalues

The proof of Theorem 3.1 resets on a cating argument based in the nature of the roots with respect to ξ_n of $p_\varphi^\pm$. At $x_n = 0$, the polynomials $p_\varphi^\pm$ in ξ_n have two complex roots. Depending on the signs of the imaginary parts of the two roots of the two poynomials, we shall adopt different strategies for the proof of partial Carleman estimates. Follow to [LR97], we set

$$\mu^\pm(x,\xi') = q_2^\pm(x,\xi') + \frac{(q_1^\pm(x,\xi'))^2}{(\partial_{x_n}\varphi^\pm)^2},$$

and we define

$$\begin{array}{lcl} \mathcal{E}^{\pm,+} & = & \{(x,\xi') \in V^+ \times \mathbb{R}^{n-1};\ \mu^\pm(x,\xi') > 0\}, \\ \mathcal{E}^{\pm,-} & = & \{(x,\xi') \in V^+ \times \mathbb{R}^{n-1};\ \mu^\pm(x,\xi') < 0\}, \\ \mathcal{Z}^\pm & = & \{(x,\xi') \in V^+ \times \mathbb{R}^{n-1};\ \mu^\pm(x,\xi') = 0\}. \end{array}$$

Remark 3.1 It is important for the sequel of the analysis to note that from the property (3.2) that the regions $\mathcal{E}^{\pm,+}$ and $\mathcal{Z}^\pm$ are bounded.

The following lemma give us an idea about the postion of the roots of $p_\varphi^\pm$ (or the eigenvalues of p_φ) in the complex plane depending to the sign of $\mu^\pm$.

Lemma 3.1 *[LR97, Lemma 3]*

1) *In the region $\mathcal{E}^{\pm,+}$, the polynomials $p_\varphi^\pm$ have two distinct roots $\rho^{\pm,+}$ and $\rho^{\pm,-}$ that satisfy $\mathrm{Im}(\rho^{\pm,+}) > 0$ and $\mathrm{Im}(\rho^{\pm,-}) < 0$.*
2) *In the region $\mathcal{E}^{\pm,-}$, the imaginary parts of the two roots have the same sign as $-(\partial_{x_n}\varphi^\pm)$.*
3) *In the region $\mathcal{Z}^\pm$, one of the roots is real.*

Remarks 3.2 1) For $|\xi'| \gg 1$, there are two roots $\rho^{+,+}$ and $\rho^{+,-}$ (resp. $\rho^{-,+}$ and $\rho^{-,-}$) for p_φ^+ (resp. p_φ^-) with $\mathrm{Im}(\rho^{+,+}) > 0$ and $\mathrm{Im}(\rho^{+,-}) < 0$ (resp. $\mathrm{Im}(\rho^{-,+}) > 0$ and $\mathrm{Im}(\rho^{-,-}) < 0$). And as the value of μ^+ (resp. μ^-) deceases, the roots $\rho^{+,+}$ (resp. $\rho^{-,-}$) move towards to real axis, and accrosse it in the region $\mathcal{Z}^+$ (resp. $\mathcal{Z}^-$).

2) Note that from the proof of [LR97, Lemma 3], we can deduce that $\mu^\pm \geq C > 0$ is equivalent to have $\mathrm{Im}(\rho^{\pm,+}) \geq C'$ and $\mathrm{Im}(\rho^{\pm,-}) \leq -C'$.

3) Morever, the roots $\rho^{\pm,+}$ and $\rho^{\pm,-}$ are in S_t^1 in fact

$$\left\{ \begin{array}{lll} \rho^{\pm,+} + \rho^{\pm,-} & = -2i(\partial_{x_n}\varphi^\pm) & \in S_t^0 \\ \rho^{\pm,+}.\rho^{\pm,-} & = q_2^\pm + 2iq_1^\pm & \in S_t^2 \end{array} \right.$$

and a simple discussion give the result.

Proposition 3.1 *With the assumptions made to the weight function φ we have*

1) $\mathcal{E}^{+,+} \subset \mathcal{E}^{-,+}$.
2) $\mathrm{d}(\mathcal{E}^{+,+}, \mathcal{Z}^-) \geq C > 0$.

This result gives meaning to the assumptions on the weight function in fact the proposition give us informations about the position of the roots each relative to other depending only to the sign of μ^+. Precisely, the root $\rho^{+,+}$ cross the real axis before the root $\rho^{-,-}$ does, as μ^+ decreases from positive to negative values. This configration will make sens in the proof of Carleman estimate since the Calderòn projector technique can not be used in situation while $\mathrm{Im}(\rho^{-,\pm}) > 0$ and the root $\rho^{+,+}$ cross the real axis.

The same result as Proposition 3.1 have been prooved in [RR10] for the operator $\partial_{x_0} - \nabla.c(x)\nabla$ where the scalar coefficient $c(x)$ is price wise smooth yet discontinous accross the interface, however the price is to make more assumptions in the weight function φ and this is according to the sign of the difference between $c(x_0, x', x_n = 0^+)$ and $c(x_0, x', x_n = 0^-)$.

Proof :

The result is clear in the case when $|\xi'| > R$ from Remark 3.1. Now for $(x,\xi') \in V^+ \times \{|\xi'| < R\}$ we have from the expression of $q_1^{\pm}$ and $q_2^{\pm}$ in (3.9) that

$$\begin{aligned}(\mu^- - \mu^+)(x,\xi')_{|x_n=0} = &\ \left((\partial_{x_n}\varphi^+)^2 - (\partial_{x_n}\varphi^-)^2\right)_{|x_n=0} \\ + &\ \left(\frac{\tilde{r}(x,\xi',\partial_{x'}\varphi^-)^2}{(\partial_{x_n}\varphi^-)^2} - \frac{\tilde{r}(x,\xi',\partial_{x'}\varphi^+)^2}{(\partial_{x_n}\varphi^+)^2}\right)_{|x_n=0} - 1 \\ = &\left(\left((\partial_{x_n}\varphi^+)^2 - (\partial_{x_n}\varphi^-)^2\right)\left(1 + \frac{\tilde{r}(x,\xi',\partial_{x'}\varphi)^2}{(\partial_{x_n}\varphi^+)^2(\partial_{x_n}\varphi^-)^2}\right)\right)_{|x_n=0} - 1 \geq C\end{aligned}$$

which come from the continuity of the weight function near the interface and the two first assumptions in (3.6). And this achieve the proof. ■

3.3 Estimation in the region $\mathcal{E}^{+,+}$

In this section we study the elliptic region $\mathcal{E}^{+,+}$ corresponding to the roots with strictly negative imaginary part for $\rho^{+,-}$ and $\rho^{-,-}$ and with strictly positive imaginary part for $\rho^{-,+}$ and $\rho^{+,+}$. Making use of the Calderòn projector technique we shall prove the following partial Carleman estimate.

Proposition 3.2 *Let $\chi^+(x,\xi') \in S_t^0$ with a compact support in x contained in V, such that* $\mathrm{supp}(\chi^+) \subset \{\mu^+(x,\xi') \geq c_0 > 0\}$. *Then there exist $C > 0$ and $h_0 > 0$ such that*

$$\begin{aligned}\|\mathrm{op}_t(\chi^+)v\|_1^2 + h|\mathrm{op}_t(\chi^+)v_{|x_n=0}|_1^2 + h|\mathrm{op}_t(\chi^+)(D_{x_n}v)_{|x_n=0}|_0^2 \leq \\ C(\|P_\varphi v\|_0^2 + h^2\|v\|_1^2 + h^4|D_{x_n}v_{|x_n=0}|_0^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2)\end{aligned}$$

for $0 < h \leq h_0$ and $v = {}^t(v^+, v^-)$; v^+, $v^- \in \mathscr{C}_c^\infty(K^+)$ and satisfying (3.7).

Proof :
We begin by noting that by Remarks 3.2 we have

$$\begin{cases}\mathrm{Im}(\rho^{\pm,+}) \geq C \text{ and } \mathrm{Im}(\rho^{\pm,-}) \leq -C \text{ in } \mathrm{supp}(\chi^+) \\ \chi^+\rho^{\pm,+},\ \chi^+\rho^{\pm,-} \in S_t^1.\end{cases}$$

We set $\begin{cases} w = \mathrm{op}_t(\chi^+)v \\ g = \mathrm{op}_t(\chi^+)P_\varphi v + \overbrace{[P_\varphi, \mathrm{op}_t(\chi^+)]}^{\in h\Psi^1}\, v\end{cases}$ which gives $P_\varphi w = g$ and

$$\|g\|_0 \leq C(\|P_\varphi v\|_0 + h\|v\|_1). \tag{3.10}$$

The transmission conditions satisfied by w are

$$\begin{cases} w^+_{|x_n=0} = w^-_{|x_n=0} + \theta_{\varphi,\chi^+} \\ (D_{x_n} + i(\partial_{x_n}\varphi^+))w^+_{|x_n=0} + (D_{x_n} + i(\partial_{x_n}\varphi^-))w^-_{|x_n=0} = G_1\end{cases} \tag{3.11}$$

where $\theta_{\varphi,\chi^+} = \mathrm{op}_t(\chi^+)\theta_{\varphi\,|x_n=0}$ and

$$\begin{aligned}G_1 = \overbrace{[(D_{x_n} + i(\partial_{x_n}\varphi^+)), \mathrm{op}_t(\chi^+)]}^{\in h\Psi_t^0}\, v^+_{|x_n=0} &+ \overbrace{[(D_{x_n} + i(\partial_{x_n}\varphi^-)), \mathrm{op}_t(\chi^+)]}^{\in h\Psi_t^0}\, v^-_{|x_n=0} \\ &+ \mathrm{op}_t(\chi^+)\tilde{\theta}_{\varphi\,|x_n=0}\end{aligned}$$

that satisfies

$$|G_1|_0 \leq C(h|v_{|x_n=0}|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.12}$$

Estimation of $\|w\|_1$: We denote by $\underline{\psi}$ the zero-extension of a function $\psi \in \mathscr{C}^\infty(V^+)$ to $\mathbb{R}^n$, $\gamma_0(\psi) = \psi_{|x_n=0}$, $\gamma_1(\psi) = D_{x_n}\psi_{|x_n=0}$ and $\delta^{(j)} = \partial^j_{x_n}(\delta_{x_n=0})$ then we have by trace formula

$$P_\varphi \underbrace{\begin{pmatrix}\gamma_0(w^+)\\ \gamma_0(w^-)\end{pmatrix}}_{\gamma_0(w)} = \underbrace{\begin{pmatrix}\underline{g}^+\\ \underline{g}^-\end{pmatrix}}_{\underline{g}} - h^2 \underbrace{\begin{pmatrix}\gamma_0(w^+)\\ \gamma_0(w^-)\end{pmatrix}}_{\gamma_0(w)} \otimes \delta' +$$

$$\frac{h}{i}\left(\underbrace{\begin{pmatrix}\gamma_1(w^+)\\ \gamma_1(w^-)\end{pmatrix}}_{\gamma_1(w)} - \underbrace{\begin{pmatrix}-2i(\partial_{x_n}\varphi^+) & 0\\ 0 & -2i(\partial_{x_n}\varphi^-)\end{pmatrix}}_{\mathrm{op}_t(q_\varphi)} \underbrace{\begin{pmatrix}\gamma_0(w^+)\\ \gamma_0(w^-)\end{pmatrix}}_{\gamma_0(w)} \right) \otimes \delta.$$

Setting

$$w_1 = \gamma_0(w) \quad \text{and} \quad w_0 = \gamma_1(w) - \mathrm{op}_t(q_\varphi)\gamma_0(w) \tag{3.13}$$

then we get

$$P_\varphi \underline{w} = \underline{g} - h^2 w_1 \otimes \delta' + \frac{h}{i} w_0 \otimes \delta.$$

Let $\chi \in S^0$ satisfying the following assumptions

$$\begin{cases} \chi \equiv 1 \text{ for } \{|\xi| \gg 1\} \text{ and in a neighborhood of } \mathrm{supp}(\chi^+) \\ \mathrm{supp}(\chi) \cap (\det(p_\varphi))^{-1}(\{0\}) = \emptyset. \end{cases}$$

These conditions are compatible in fact $\det(p_\varphi)(x,\xi) = 0$ is equivalent to $p^+_\varphi(x,\xi) = 0$ or $p^-_\varphi(x,\xi) = 0$ and this mean that $(x,\xi') \in \mathcal{Z}^+ \cup \mathcal{Z}^-$, however $\mathrm{d}(\mathcal{E}^{+,+}, \mathcal{Z}^\pm) \geq 0$ then our choice is justify.

From the ellipticity of p_φ on $\mathrm{supp}(\chi)$, for large M there exists a PDO, $E_M = \mathrm{Op}(e)$ with $e \in S^{-2}$ of the form $e = \Sigma^M_{j=0} h^j e_j$, with $e_j \in S^{-2-j}$ and $e^\pm_0 = \frac{\chi}{p^\pm_\varphi}$, that satisfies

$$E_M \circ P_\varphi = \mathrm{Op}(\chi) + h^{m+1} R_M, \quad R_M \in \Psi^{-M-1}.$$

Note that the symbols $e^\pm_j$, $j = 1, \ldots, M$ are in the form of rational functions for large $|\xi_n|$ with $\rho^{\pm,+}$ and $\rho^{\pm,-}$ for only poles. Then we obtain

$$\underline{w} = E_M \underline{g} + E_M(-h^2 w_1 \otimes \delta' + \frac{h}{i} w_0 \otimes \delta) + g_1$$

where $g_1 = \mathrm{Op}(1-\chi)(\underline{w}) - h^{M+1} R_M \underline{w}$. With Lemma A.1 we have $\mathrm{Op}(1-\chi) \circ \mathrm{op}_t(\chi^+) \in \bigcap_{N\in\mathbb{N}} h^N \Psi^{-N}$ which imply

$$\|g_1\|_2 \leq C h^2 \|v\|_0. \tag{3.14}$$

For $j = 0, 1$ we set

$$\tilde{t}_j(x,\xi') = \frac{1}{2i\pi} \int \mathrm{e}^{\frac{i}{h} x_n \xi_n} e(x,\xi',\xi_n) \xi^j_n \,\mathrm{d}\xi_n$$

then we have

$$E_M(\frac{h}{i} w_0 \otimes \delta)(x) = \frac{1}{(2\pi h)^n} \iint \mathrm{e}^{\frac{i}{h}\langle x'-y',\xi'\rangle} \tilde{t}_0(x,\xi') w_0(y') \,\mathrm{d}y' \,\mathrm{d}\xi'$$
$$E_M(-h^2 w_1 \otimes \delta')(x) = \frac{1}{(2\pi h)^n} \iint \mathrm{e}^{\frac{i}{h}\langle x'-y',\xi'\rangle} \tilde{t}_1(x,\xi') w_1(y') \,\mathrm{d}y' \,\mathrm{d}\xi'.$$

Note that the integral defining $\tilde{t}_0$ is absolutly converging but whos defining $\tilde{t}_1$ must be understood in the sens of oscillatry integrals [Hör85, Mar02] and we can write it in the form

$$\tilde{t}^\pm_1(x,\xi') = \frac{1}{2i\pi} D_{y_n} \left(\int \mathrm{e}^{\frac{i}{h} y_n \xi_n} e^\pm(x,\xi',\xi_n) \,\mathrm{d}\xi_n \right)_{|y_n = x_n}.$$

With the choise of the cut-off function χ we can see that $e(x,\xi)$ is holomorphic for large $|\xi_n|$, $\xi_n \in \mathbb{C}$, thus we obtain for $x_n > 0$

$$\tilde{t}_0^{\pm}(x,\xi',h) = \frac{1}{2i\pi}\int_{\gamma^{\pm}} \mathrm{e}^{\frac{i}{h}x_n\xi_n} e^{\pm}(x,\xi',\xi_n)\,\mathrm{d}\xi_n \tag{3.15}$$

$$\tilde{t}_1^{\pm}(x,\xi',h) = \frac{1}{2i\pi}\int_{\gamma^{\pm}} \mathrm{e}^{\frac{i}{h}x_n\xi_n} e^{\pm}(x,\xi',\xi_n)\xi_n\,\mathrm{d}\xi_n \tag{3.16}$$

where $\gamma^{\pm}$ the union of the segment $\{\xi_n \in \mathbb{R}^n;\ |\xi_n| \leq C_0|\xi'|\}$ and the half circle $\{\xi_n \in \mathbb{C};\ |\xi_n| = C_0|\xi'|,\ \mathrm{Im}(\xi_n) > 0\}$ where C_0 is chosen sufficiently large so as to have the roots $\rho^{\pm,+}$ inside the domain with boundary $\gamma^{\pm}$. The exprssions (3.15) and (3.16) are valide also for $x_n = 0$ and for $j = 0,1$ we have

$$|D_{x_n}^l \partial_{x'}^{\alpha} \partial_{\xi'}^{\beta} \tilde{t}_j^{\pm}(x,\xi',h)| \leq C_{\alpha,\beta,l}\langle \xi' \rangle^{-1+j+l-|\beta|} \quad x_n \geq 0,\ l \in \mathbb{N}. \tag{3.17}$$

Let $\chi_1(x,\xi') \in S_t^0$ satisfying the same requirement as χ^+, such that

$$\begin{cases} \chi_1(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi^+) \\ \chi(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi_1). \end{cases}$$

We set $t_j = \chi_1 \tilde{t}_j$; $j = 0,1$ and $g_2 = \mathrm{op}_t((1-\chi_1)\tilde{t}_0)w_0 + \mathrm{op}_t((1-\chi_1)\tilde{t}_1)w_1$ and we write

$$\underline{w} = E_M \underline{g} + \mathrm{op}_t(t_0)w_0 + \mathrm{op}_t(t_1)w_1 + g_1 + g_2. \tag{3.18}$$

Now from (3.17) if we show t_j as a symbol in $\mathbb{R}^{n-1}$ with x_n as a parameter, then while $\mathrm{supp}(1-\chi_1) \cap \mathrm{supp}(\chi^+) = \emptyset$ then by composition formula and Remark A.1 we obtain

$$\|g_2\|_2 \leq Ch^3(|v_{|x_n=0}|_{-1} + |(D_{x_n}v)_{|x_n=0}|_{-1}) \leq Ch^2(\|v\|_1 + |(D_{x_n}v)_{|x_n=0}|_0). \tag{3.19}$$

While $e(x,\xi)$ is holomorphic relative to ξ_n in $\mathrm{supp}(\chi_1)$, we can then write for $j = 0,1$ that

$$t_j^{\pm}(x,\xi',h) = \chi_1(x,\xi')\frac{1}{2i\pi}\int_{\gamma_0^{\pm}} e^{\frac{i}{h}x_n\xi_n} e^{\pm}(x,\xi',\xi_n)\xi_n^j\,\mathrm{d}\xi_n \tag{3.20}$$

where $\gamma_0^{\pm}$ is a direct contour surrounding the roots $\rho^{\pm,+}$ in the region $\mathrm{Im}(\xi_n) \geq C_0|\xi'|$ for $C_0 > 0$. In $\mathrm{supp}(\chi_1)$ we have

$$e_0^{\pm} = \frac{1}{p_{\varphi}^{\pm}} = \frac{1}{\rho^{\pm,+} - \rho^{\pm,-}}\left(\frac{1}{\xi_n - \rho^{\pm,+}} - \frac{1}{\xi_n - \rho^{\pm,-}}\right).$$

Then the residue formula yields for $j = 0,1$ that

$$\mathrm{e}^{-\frac{i}{h}x_n\rho^{\pm,+}} t_j^{\pm}(x,\xi',h) = \chi_1(x,\xi')\frac{(\rho^{\pm,+})^j}{\rho^{\pm,+} - \rho^{\pm,-}} + h\lambda_j a^{\pm}, \quad \lambda_j^{\pm} \in S_t^{j-2}.$$

From (3.20) we get for $j = 0,1$ that

$$|D_{x_n}^l \partial_{x'}^{\alpha} \partial_{\xi'}^{\beta} t_j^{\pm}(x,\xi',h)| \leq C_{\alpha,\beta,l}\mathrm{e}^{-\frac{C_0}{h}x_n\langle\xi'\rangle}\langle \xi' \rangle^{-1+j+l-|\beta|} \quad x_n \geq 0,\ l \in \mathbb{N}.$$

Which yields

$$|\mathrm{e}^{\frac{C_0}{h}x_n} D_{x_n}^l t_j^{\pm}(x,\xi',h)| \leq C_l \mathrm{e}^{\frac{C_0}{h}x_n(1-\langle\xi'\rangle)}\langle\xi'\rangle^{j-1+l} \leq C_l\langle\xi'\rangle^{j-1+l}.$$

It follows that

$$\begin{cases} \|\Lambda_t^1 \circ \mathrm{op}_t(t_j)w_j\|_0^2 = \displaystyle\int_{x_n>0} \mathrm{e}^{-\frac{2}{h}C_0x_n}|\mathrm{op}_t(\mathrm{e}^{\frac{C_0}{h}x_n}t_j)w_j|_1^2(x_n)\,\mathrm{d}x_n \leq Ch|w_j|_j^2 \\ \|D_{x_n} \circ \mathrm{op}_t(t_j)w_j\|_0^2 = \displaystyle\int_{x_n>0} \mathrm{e}^{-\frac{2}{h}C_0x_n}|\mathrm{op}_t(\mathrm{e}^{\frac{C_0}{h}x_n}D_{x_n}t_j)w_j|_1^2(x_n)\,\mathrm{d}x_n \leq Ch|w_j|_j^2, \end{cases}$$

which can be writen as follows

$$\|\mathrm{op}_t(t_j)w_j\|_1^2 \le Ch|w_j|_j^2. \tag{3.21}$$

From (3.18) and estimates (3.19), (3.10), (3.14) and (3.21) we obtain

$$\|w\|_1 \le C(\|P_\varphi v\|_0 + h\|v\|_1 + h^{\frac{1}{2}}(|w_0|_0 + |w_1|_1) + h^2|(D_{x_n}v)_{|x_n=0}|_0). \tag{3.22}$$

Estimation of the boundary terms $|w_0|_0$ and $|w_1|_1$: We take the trace at $x_n = 0$ of (3.18) which gives

$$\gamma_0(w) = \mathrm{op}_t(a)\gamma_0(w) + \mathrm{op}_t(b)\gamma_1(w) + G_2, \tag{3.23}$$

where $a \in S_t^0$ and $b \in S_t^{-1}$ with principal symbols

$$\left\{\begin{array}{l} a_0 = \begin{pmatrix} a_0^+ & 0 \\ 0 & a_0^- \end{pmatrix}; \quad a_0^\pm = -\left(\chi_1^\pm \dfrac{\rho^{\pm,-}}{\rho^{\pm,+} - \rho^{\pm,-}}\right)_{|x_n=0} \\ b_{-1} = \begin{pmatrix} b_{-1}^+ & 0 \\ 0 & b_{-1}^- \end{pmatrix}; \quad b_{-1}^\pm = \left(\chi_1^\pm \dfrac{1}{\rho^{\pm,+} - \rho^{\pm,-}}\right)_{|x_n=0} \end{array}\right.$$

and $G_2 = (E_M\underline{g} + g_1 + g_2)_{|x_n=0}$. Which give by Remark A.1 and estimates (3.19), (3.10) and (3.14)

$$|G_2|_1 \le Ch^{-\frac{1}{2}}\|E_M g + g_1 + g_2\|_2 \le Ch^{-\frac{1}{2}}(\|P_\varphi v\|_0 + h\|v\|_1 + h^2|(D_{x_n}v)_{|x_n=0}|_0). \tag{3.24}$$

The transmission conditions (3.11) gives

$$\left\{\begin{array}{l} \gamma_0(w^-) = \gamma_0(w^+) - \theta_{\varphi,\chi^+} \\ \gamma_1(w^-) = -\gamma_1(w^+) + k\gamma_0(w_0^+) + \widetilde{G}_1 \end{array}\right. \tag{3.25}$$

where $k = -i((\partial_{x_n}\varphi^-)_{|x_n=0} + (\partial_{x_n}\varphi^+)_{|x_n=0})$ and $\widetilde{G}_1 = i(\partial_{x_n}\varphi^-)_{|x_n=0}\theta_{\varphi,\chi^+} + G_1$, in particular we have by Remark A.1 and (3.12) that

$$|\widetilde{G}_1|_0 \le C(h|v_{|x_n=0}|_0 + |\theta_\varphi|_0 + |\tilde{\theta}_\varphi|_0) \le C(h^{\frac{1}{2}}\|v\|_1 + |\theta_\varphi|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.26}$$

From (3.23) and (3.25) we obtain

$$\mathrm{op}_t(1-a)\begin{pmatrix} \gamma_0(w^+) \\ \gamma_0(w^+) - \theta_{\varphi,\chi^+} \end{pmatrix} = \mathrm{op}_t(b)\begin{pmatrix} \gamma_1(w^+) \\ -\gamma_1(w^+) + k\gamma_0(w^+) + \widetilde{G}_1 \end{pmatrix} + G_2.$$

Thus we have

$$\begin{pmatrix} \mathrm{op}_t(1-a^+) & -\mathrm{op}_t(b^+) \\ \mathrm{op}_t(1-a^-) - \mathrm{op}_t(b^-)\circ k & \mathrm{op}_t(b^-) \end{pmatrix}\begin{pmatrix} \gamma_0(w^+) \\ \gamma_1(w^+) \end{pmatrix} =$$
$$\mathrm{op}_t(1-a)\begin{pmatrix} 0 \\ \theta_{\varphi,\chi^+} \end{pmatrix} + \mathrm{op}_t(b)\begin{pmatrix} 0 \\ \widetilde{G}_1 \end{pmatrix} + G_2$$

Thus we obtain a system of the form

$$\mathrm{op}_t(\kappa)\begin{pmatrix} \gamma_0(w^+) \\ \Lambda_t^{-1}\gamma_1(w^+) \end{pmatrix} = G_2 + \mathrm{op}_t(\sigma)\theta_{\varphi,\chi^+} + \mathrm{op}_t(\Sigma)\widetilde{G}_1,$$

where κ is a 2×2 matrix with entries in S_t^0 and σ and Σ are 2×1 matrix with entries in S_t^0 and S_t^{-1} respectively, with principal symbols

$$\kappa_0 = \begin{pmatrix} 1 - a_0^+ & -b_{-1}^+\langle\xi'\rangle \\ (1-a_0^-) - kb_{-1}^- & b_{-1}^-\langle\xi'\rangle \end{pmatrix}, \quad \sigma_0 = \begin{pmatrix} 0 \\ 1 - a_0^- \end{pmatrix}, \quad \Sigma_{-1} = \begin{pmatrix} 0 \\ b_{-1}^- \end{pmatrix}.$$

Choosing $\chi_2(x,\xi) \in S_t^0$, satisfing the same requirement as χ^+, such that

$$\begin{cases} \chi_2(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi^+) \\ \chi_1(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi_2). \end{cases}$$

In supp(χ_2) we have

$$\kappa_{0\,|\mathrm{supp}(\chi_2)} = \begin{pmatrix} \dfrac{\rho^{+,+}}{\rho^{+,+}-\rho^{+,-}} & -\langle\xi'\rangle\dfrac{1}{\rho^{+,+}-\rho^{+,-}} \\ \dfrac{\rho^{-,+}-k}{\rho^{-,+}-\rho^{-,-}} & \langle\xi'\rangle\dfrac{1}{\rho^{-,+}-\rho^{-,-}} \end{pmatrix}_{|x_n=0}$$

Since $\mathrm{Im}(\rho^{-,+}+\rho^{+,+}) \geq C\langle\xi'\rangle$ in supp(χ_2) and $\mathrm{Im}(-k) = ((\partial_{x_n}\varphi^-)+(\partial_{x_n}\varphi^+))_{|x_n=0} > 0$ by (3.6) then we find

$$|\det(\kappa_0)_{|\mathrm{supp}(\chi_2)}| = \left|\frac{\langle\xi'\rangle(\rho^{-,+}+\rho^{+,+}-k)}{(\rho^{+,+}-\rho^{+,-})(\rho^{-,+}-\rho^{-,-})}\right|_{x_n=0} \geq C.$$

It follows that κ is elliptic in supp(χ_2). Then there exists $l_M \in S_t^0$, such that

$$\mathrm{op}_t(l_M)\circ\mathrm{op}_t(\kappa) = \mathrm{op}_t(\chi_2) + h^{M+1}\widetilde{R}_M; \quad \widetilde{R}_M \in \Psi^{-M-1}$$

for $M \in \mathbb{N}$ large. This yields

$$\begin{aligned} \begin{pmatrix} \gamma_0(w^+) \\ \Lambda_t^{-1}\gamma_1(w^+) \end{pmatrix} &= \mathrm{op}_t(l_M)G_2 + \mathrm{op}_t(l_M)\circ\mathrm{op}_t(\sigma)\theta_{\varphi,\chi^+} + \mathrm{op}_t(l_M)\circ\mathrm{op}_t(\Sigma)\widetilde{G}_1 \\ &+ \;(\mathrm{op}_t(1-\chi_2)+h^{M+1}\widetilde{R}_M)\begin{pmatrix} \gamma_0(w^+) \\ \Lambda_t^{-1}\gamma_1(w^+) \end{pmatrix}. \end{aligned}$$

Since $\mathrm{supp}(1-\chi_2)\cap\mathrm{supp}(\chi^+) = \emptyset$ then by Remark A.1 and estimates (3.24) and (3.26) we obtain

$$\begin{aligned} |\gamma_0(w^+)|_1 + |\gamma_1(w^+)|_0 &\leq C(|G_2|_1 + |\theta_{\varphi,\chi^+}|_1 + |\widetilde{G}_1|_0 + h^2|v_{|x_n=0}|_0 + h^2|(D_{x_n}v)_{|x_n=0}|_0) \\ &\leq C(h^{\frac{-1}{2}}\|P_\varphi v\|_0 + h^{\frac{1}{2}}\|v\|_1 + h^{\frac{3}{2}}|(D_{x_n}v)_{|x_n=0}|_0 + |\theta_\varphi|_1 + |\tilde{\theta}_\varphi|_0). \end{aligned} \tag{3.27}$$

From the transmission conditions (3.25) and estimate (3.26) we have

$$\begin{aligned} &|\gamma_0(w^-)|_1 \leq C(h|\gamma_0(w^+)|_1 + |\theta_\varphi|_1) \\ &|\gamma_1(w^-)|_0 \leq C(|\gamma_1(w^+)|_0 + |\gamma_0(w^+)|_1 + h^{\frac{1}{2}}\|v\|_1 + |\theta_\varphi|_1 + |\tilde{\theta}_\varphi|_0). \end{aligned}$$

By summing these two last estimates we get by (3.27) that

$$|\gamma_0(w^-)|_1 + |\gamma_1(w^-)|_0 \leq C(h^{\frac{-1}{2}}\|P_\varphi v\|_0 + h^{\frac{1}{2}}\|v\|_1 + h^{\frac{3}{2}}|(D_{x_n}v)_{|x_n=0}|_0 + |\theta_\varphi|_1 + |\tilde{\theta}_\varphi|_0). \tag{3.28}$$

Then by (3.13) and estimates (3.27) and (3.28) we obtain

$$h^{\frac{1}{2}}(|w_0|_0 + |w_1|_1) \leq C(\|P_\varphi v\|_0 + h\|v\|_1 + h^2|(D_{x_n}v)_{|x_n=0}|_0 + h^{\frac{1}{2}}|\theta_\varphi|_1 + h^{\frac{1}{2}}|\tilde{\theta}_\varphi|_0). \tag{3.29}$$

End of proof: Observing that

$$D_{x_n}\mathrm{op}_t(\chi^+)v = \mathrm{op}_t(\chi^+)(D_{x_n}v) + \underbrace{[D_{x_n},\mathrm{op}_t(\chi^+)]}_{\in h\Psi_t^0}v.$$

Then by (3.27), (3.28) and Remark A.1 we obtain

$$\begin{aligned} h^{\frac{1}{2}}(|\mathrm{op}_t(\chi^+)v_{|x_n=0}|_1 + |\mathrm{op}_t(\chi^+)(D_{x_n}v)_{|x_n=0}|_0) &\leq Ch^{\frac{1}{2}}(|w_{|x_n=0}|_1 + |D_{x_n}w_{|x_n=0}|_0 + h|v_{|x_n=0}|_0) \\ &\leq C(\|P_\varphi v\|_0 + h\|v\|_1 + h^2|(D_{x_n}v)_{|x_n=0}|_0 + h^{\frac{1}{2}}|\theta_\varphi|_1 + h^{\frac{1}{2}}|\tilde{\theta}_\varphi|_0). \end{aligned} \tag{3.30}$$

Thus we conclude the proof by combining estimates (3.22), (3.29) and (3.30). ■

3.4 Estimation in the region $\mathcal{E}^{+,-}$

With a microlocal cutt-off function, we place ourselves in the region $\mathcal{E}^{+,-}$, finitly away from $\mathcal{Z}^+$. Making use of the standard techniques to prove Carleman estimates for both P_φ^- and P_φ^+.

Proposition 3.3 *Let $\chi^-(x,\xi') \in S_t^0$ with a compact support in x contained in V, such that* $\operatorname{supp}(\chi^-) \subset \{\mu^+(x,\xi') \le -C < 0\}$. *Then there exist $C > 0$ and $h_0 > 0$ such that*

$$h\|\operatorname{op}_t(\chi^-)v\|_1^2 + h|\operatorname{op}_t(\chi^-)v_{|x_n=0}|_1^2 + h|\operatorname{op}_t(\chi^-)(D_{x_n}v)_{|x_n=0}|_0^2 \le C(\|P_\varphi v\|_0^2 + h^2\|v\|_1^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2)$$

for $0 < h \le h_0$ and $v = {}^t(v^+, v^-)$; v^+, $v^- \in \mathscr{C}_c^\infty(K^+)$ that satisfy (3.7).

Proof :
We set $w = \operatorname{op}_t(\chi^-)v$ and $g = \operatorname{op}_t(\chi^-)P_\varphi v + [P_\varphi, \operatorname{op}_t(\chi^-)]v$ then $P_\varphi w = g$ and we have

$$\|g\|_0 \le C(\|P_\varphi v\|_0 + h\|v\|_1). \tag{3.31}$$

As in the previous proof the transmission conditions satisfyed by w^- and w^+ are

$$\begin{cases} w^+_{|x_n=0} = w^-_{|x_n=0} + \theta_{\varphi,\chi^-} \\ (D_{x_n} + i(\partial_{x_n}\varphi^+))w^+_{|x_n=0} + (D_{x_n} + i(\partial_{x_n}\varphi^-))w^-_{|x_n=0} = G_1 \end{cases} \tag{3.32}$$

where $\theta_{\varphi,\chi^-} = \operatorname{op}_t(\chi^-)\theta_{\varphi\,|x_n=0}$ and

$$\begin{aligned} G_1 = [(D_{x_n} + i(\partial_{x_n}\varphi^+)), \operatorname{op}_t(\chi^-)]v^+_{|x_n=0} &+ [(D_{x_n} + i(\partial_{x_n}\varphi^-)), \operatorname{op}_t(\chi^-)]v^-_{|x_n=0} \\ &+ \operatorname{op}_t(\chi^-)\tilde{\theta}_{\varphi\,|x_n=0} \end{aligned}$$

that satisfies

$$|G_1|_0 \le C(h|v_{|x_n=0}|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.33}$$

From [LR95, Lemme 2] we have

$$h\|w^\pm\|_1^2 + \operatorname{Re}\left(h\mathcal{B}^\pm(w^\pm) + h^2\left((D_{x_n}w^\pm + L_1^\pm w^\pm)_{|x_n=0}, L_0^\pm w^\pm_{|x_n=0}\right)_0\right) \le C\|g^\pm\|_0^2 \tag{3.34}$$

for h sufficiently small, where $L_1^\pm \in \mathcal{D}_t^1$, $L_0^\pm \in \Psi_t^0$ and $\mathcal{B}^\pm$ are the quadratic forms given by

$$\mathcal{B}^\pm(\psi) = \left(\overbrace{\begin{pmatrix} 2(\partial_{x_n}\varphi^\pm) & B_1^\pm\Lambda_t^{-1} \\ \Lambda_t^{-1}B_1^{\pm'} & \Lambda_t^{-1}B_2^\pm\Lambda_t^{-1} \end{pmatrix}}^{=B^\pm\in\Psi_t^0} \begin{pmatrix} \gamma_1(\psi) \\ \Lambda_t^1\gamma_0(\psi) \end{pmatrix}, \begin{pmatrix} \gamma_1(\psi) \\ \Lambda_t^1\gamma_0(\psi) \end{pmatrix}\right)_0 \tag{3.35}$$

where $B_1^\pm$, $B_1^{\pm'} \in \mathcal{D}_t^1$ with $\sigma(B_1^\pm) = \sigma(B_1^{\pm'}) = 2q_1^\pm$ and $B_2^\pm \in \mathcal{D}_t^2$ with $\sigma(B_2^\pm) = -2(\partial_{x_n}\varphi^\pm)q_2^\pm$. In particular, we have

$$h^2\left|\left((D_{x_n}w^\pm + L_1^\pm w^\pm)_{|x_n=0}, L_0^\pm w^\pm_{|x_n=0}\right)_0\right| \le Ch^2(|\gamma_1(w^\pm)|_0^2 + |\gamma_0(w^\pm)|_1^2). \tag{3.36}$$

The principal symbols of $B^\pm$ are

$$\sigma(B^\pm) = \begin{pmatrix} 2(\partial_{x_n}\varphi^\pm) & 2q_1^\pm\langle\xi'\rangle^{-1} \\ 2q_1^\pm\langle\xi'\rangle^{-1} & -2(\partial_{x_n}\varphi^\pm)q_2^\pm\langle\xi'\rangle^{-2} \end{pmatrix}. \tag{3.37}$$

We observe that $\det(\sigma(B^\pm)) = -4(\partial_{x_n}\varphi^\pm)^2\langle\xi'\rangle^{-2}\mu^\pm$, it follows that in $\operatorname{supp}(\chi^-)$ we have that $\det(\sigma(B^+)) \ge C$. Morever since $(\partial_{x_n}\varphi^+)_{|x_n=0} > 0$ it folows that $\sigma(B^+)$ is positive defined. The transmission condition (3.32) can be written as follows

$$\begin{pmatrix} \gamma_1(w^-) \\ \Lambda_t^1\gamma_0(w^-) \end{pmatrix} = \underbrace{\begin{pmatrix} -1 & k\Lambda_t^{-1} \\ 0 & h \end{pmatrix}}_{C^-} \begin{pmatrix} \gamma_1(w^+) \\ \Lambda_t^1\gamma_0(w^+) \end{pmatrix} + \begin{pmatrix} \widetilde{G}_1 \\ -\Lambda_t^1\theta_{\varphi,\chi^-} \end{pmatrix} \tag{3.38}$$

where $k = -i(h(\partial_{x_n}\varphi^-)_{|x_n=0} + (\partial_{x_n}\varphi^+)_{|x_n=0})$ and $\widetilde{G}_1 = i(\partial_{x_n}\varphi^-)_{|x_n=0}\theta_{\varphi,\chi^-} + G_1$ that satisfies from (3.33) and Remark A.1

$$|\widetilde{G}_1|_0 \leq C(h^{\frac{1}{2}}\|v\|_1 + |\theta_\varphi|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.39}$$

Then we obtain by (3.35) and (3.38) that

$$\mathcal{B}^-(w^-) = \left(\widetilde{B}^-\begin{pmatrix} \gamma_1(w^+) \\ \Lambda_t^1\gamma_0(w^+)\end{pmatrix}, \begin{pmatrix} \gamma_1(w^+) \\ \Lambda_t^1\gamma_0(w^+)\end{pmatrix}\right) + U(\gamma_1(w^+), \gamma_0(w^+), \theta_{\varphi,\chi^-}, \widetilde{G}_1)$$

where $\sigma(\widetilde{B}^-) = {}^t\overline{\sigma(C^-)}\sigma(B^-)\sigma(C^-) \in S_t^0$ and

$$U(\gamma_1(w^+), \gamma_0(w^+), \theta_{\varphi,\chi^-}, \widetilde{G}_1) = \left(B^-C^-\begin{pmatrix} \gamma_1(w^+) \\ \Lambda_t^1\gamma_0(w^+)\end{pmatrix}, \begin{pmatrix} \widetilde{G}_1 \\ \Lambda_t^1\theta_{\varphi,\chi^-}\end{pmatrix}\right)_0 +$$
$$\left(B^-\begin{pmatrix} \widetilde{G}_1 \\ \Lambda_t^1\theta_{\varphi,\chi^-}\end{pmatrix}, C^-\begin{pmatrix} \gamma_1(w^+) \\ \Lambda_t^1\gamma_0(w^+)\end{pmatrix}\right)_0 + \left(B^-\begin{pmatrix} \widetilde{G}_1 \\ \Lambda_t^1\theta_{\varphi,\chi^-}\end{pmatrix}, \begin{pmatrix} \widetilde{G}_1 \\ \Lambda_t^1\theta_{\varphi,\chi^-}\end{pmatrix}\right)_0.$$

Which satisfies from (3.33), (3.39) and Remark A.1 the following estimate

$$|U(\gamma_1(w^+), \gamma_0(w^+), \theta_{\varphi,\chi^-}, \widetilde{G}_1)| \leq C(|\gamma_0(w^+)|_1^2 + |\gamma_1(w^+)|_0^2 + h\|v\|_1^2 + |\theta_\varphi|_1^2 + |\tilde{\theta}_\varphi|_0^2). \tag{3.40}$$

So that, for any $K > 0$ and for $\alpha > 0$ sufficiently large, we can suppose that $\alpha\sigma(B^+) + \sigma(\widetilde{B}^-) \geq K > 0$. Hence, by Gårding inequality (Lemma A.2) and the estimate (3.40) we obtain for all $0 < h \leq h_0$ that

$$\mathrm{Re}(\alpha\mathcal{B}^+(w^+) + \mathcal{B}^-(w^-)) \geq \frac{K}{2}(|\gamma_0(w^+)|_1^2 + |\gamma_1(w^+)|_0^2) - C(h\|v\|_1^2 + |\theta_\varphi|_1^2 + |\tilde{\theta}_\varphi|_0^2). \tag{3.41}$$

The transmission conditions (3.38) and the estimate (3.39) yields

$$|\gamma_0(w^-)|_1^2 + |\gamma_1(w^-)|_0^2 \leq C(|\gamma_0(w^+)|_1^2 + |\gamma_1(w^+)|_0^2 + h\|v\|_1^2 + |\theta_\varphi|_1^2 + |\tilde{\theta}_\varphi|_0^2). \tag{3.42}$$

A similary estimate to (3.41) can now be obtained for w^- instead of w^+ in the right hand side by estimate (3.42) then for all $0 < h \leq h_0$ we find

$$\mathrm{Re}(\alpha\mathcal{B}^+(w^+) + \mathcal{B}^-(w^-)) \geq \frac{K}{2}(|\gamma_0(w^-)|_1^2 + |\gamma_1(w^-)|_0^2) - C(h\|v\|_1^2 + |\theta_\varphi|_1^2 + |\tilde{\theta}_\varphi|_0^2). \tag{3.43}$$

With the sum of $(3.34)^-$ and $\alpha(3.34)^+$; and from estimates (3.36), (3.41) and (3.43) we obtain

$$\begin{aligned} &h|\gamma_0(w^-)|_1^2 + h|\gamma_1(w^-)|_0^2 + h|\gamma_0(w^+)|_1^2 + h|\gamma_1(w^+)|_0^2 + h\|w^-\|_1^2 + h\|w^+\|_1^2 \\ &\qquad \leq C(\|g^+\|_0^2 + \|g^-\|_0^2 + h^2\|v\|_1^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2). \end{aligned} \tag{3.44}$$

Finally with taking into account of the expression

$$D_{x_n}\mathrm{op}_t(\chi^-)v = \mathrm{op}_t(\chi^-)(D_{x_n}v) + \underbrace{[D_{x_n}, \mathrm{op}_t(\chi^-)]}_{\in h\Psi_t^0} v,$$

and estimates (3.31) and (3.44) then with proceeding as in the end of the previous section we obtain the result for h sufficiently small. ∎

3.5 Estimation in the neighborhood of the region $\mathcal{Z}^+$

With a cut-off function, we place ourselves around the region $\mathcal{Z}^+$, while staying in the region $\mathcal{E}^{-,+}$ away from $\mathcal{Z}^-$. We adopte the technique of Calderòn projector for P_φ^- and standard techniques for P_φ^+.

Lemma 3.2 *There exist $c_0 > 0$ sufficienty small and C', $C'' \geq 0$ such that*
(3.45)

$$\forall\, x \in V^+,\ \xi' \in \mathbb{R}^{n-1};\ |\mu^+(x,\xi')| \leq c_0 \Longrightarrow \begin{cases} \mu^-(x,\xi') \geq C'. \\ \Big(((\partial_{x_n}\varphi^-) + (\partial_{x_n}\varphi^+))^2 - \mu^+(x,\xi')\Big)_{|x_n=0} \geq C''. \end{cases}$$

Proof :
The first property of (3.45) follow easly from Proposition 3.1 by choosing c_0 small. For the second one we have just to choose $\epsilon > 0$ and $C'' > 0$ sufficiently small such that $((\partial_{x_n}\varphi^-)+(\partial_{x_n}\varphi^+))^2_{|x_n=0} \geq C'' + c_0$ which is compatible from the assumptions (3.6). ■

Proposition 3.4 *Let $\chi^0(x,\xi') \in S^0_t$ with a compact support in x contained in V, such that for $c_0 > 0$ sufficiently small $\{|\mu^+(x,\xi')| \leq c_0\}$ in* $\operatorname{supp}(\chi^0)$. *Then there exist $C > 0$ and $h_0 > 0$ such that*

$$\begin{aligned} h\|\mathrm{op}_t(\chi^0)v\|_1^2 + h|\mathrm{op}_t(\chi^0)v_{|x_n=0}|_1^2 + h|\mathrm{op}_t(\chi^0)(D_{x_n}v)_{|x_n=0}|_0^2 &\leq \\ C(\|P_\varphi v\|_0^2 + h^2\|v\|_1^2 + h^4|(D_{x_n}v)_{|x_n=0}|_0^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2) \end{aligned}$$

for $0 < h \leq h_0$ and $v = {}^t(v^+, v^-)$; v^+, $v^- \in \mathscr{C}^\infty_c(K^+)$ and satisfying (3.7).

Proof :
We choose $c_0 > 0$ that satisfy the conclusion of Lemma 3.2, then by Remarks 3.2 we have $\mathrm{Im}(\rho^{-,+}) \geq C$ and $\mathrm{Im}(\rho^{-,-}) \leq -C$. We set $w = \mathrm{op}_t(\chi^0)v$ and $g = \mathrm{op}_t(\chi^0)P_\varphi v + [P_\varphi, \mathrm{op}_t(\chi^0)]v$. Then we have $P_\varphi w = g$ and

$$\|g^\pm\|_0 \leq C(\|P^\pm_\varphi v^\pm\|_0 + h\|v^\pm\|_1). \tag{3.46}$$

The transmission conditions satisfied by w are

$$\begin{cases} w^+_{|x_n=0} = w^-_{|x_n=0} + \theta_{\varphi,\chi^0} \\ (D_{x_n} + i(\partial_{x_n}\varphi^+))w^+_{|x_n=0} + (D_{x_n} + i(\partial_{x_n}\varphi^-))w^-_{|x_n=0} = G_1 \end{cases} \tag{3.47}$$

where $\theta_{\varphi,\chi^0} = \mathrm{op}_t(\chi^0)\theta_{\varphi\,|x_n=0}$ and

$$\begin{aligned} G_1 = \overbrace{[(D_{x_n} + i(\partial_{x_n}\varphi^+)), \mathrm{op}_t(\chi^0)]}^{\in\, h\Psi^0_t}\, v^+_{|x_n=0} \quad &+ \quad \overbrace{[(D_{x_n} + i(\partial_{x_n}\varphi^-)), \mathrm{op}_t(\chi^0)]}^{\in\, h\Psi^0_t}\, v^-_{|x_n=0} \\ &+ \quad \mathrm{op}_t(\chi^0)\tilde{\theta}_{\varphi\,|x_n=0} \end{aligned}$$

that satisfies

$$|G_1|_0 \leq C(h|v_{|x_n=0}|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.48}$$

Calderòn projector technique applied to P^-_φ: We keep the same notations as in proof of Proposition 3.2. Thus we obtain thanks to the first property of Lemma 3.2 an estimate of the same form as (3.22), namely we obtain

$$\|w^-\|_1 \leq C(\|P^-_\varphi v^-\|_0 + h\|v^-\|_1 + h^{\frac{1}{2}}(|\gamma_1(w^-)|_0 + |\gamma_0(w^-)|_1) + h^2|(D_{x_n}v^-)_{|x_n=0}|_0). \tag{3.49}$$

Also we have the following trace formula, of the same form as (3.23), namely

$$\gamma_0(w^-) = \mathrm{op}_t(a^-)\gamma_0(w^-) + \mathrm{op}_t(b^-)\gamma_1(w^-) + G^-_2 \tag{3.50}$$

where $a^- \in S^0_t$, $b^- \in S^{-1}_t$ with principal symbols

$$a^-_0 = -\left(\chi_1 \frac{\rho^{-,-}}{\rho^{-,+} - \rho^{-,-}}\right)_{|x_n=0} \quad ; \quad b^-_{-1} = \left(\chi_1 \frac{1}{\rho^{-,+} - \rho^{-,-}}\right)_{|x_n=0} \tag{3.51}$$

where $\chi_1 \in S_t^0$, satisfying the same requirement as χ^0 and is equal to one in a neighborhood of $\mathrm{supp}(\chi^0)$. And G_2^- satisfies

$$|G_2^-|_1 \leq Ch^{-\frac{1}{2}}(\|P_\varphi^- v^-\|_0 + h\|v^-\|_1 + h^2|(D_{x_n}v^-)_{|x_n=0}|_0). \tag{3.52}$$

Then with the transmission conditions (3.47) we can write

$$\begin{cases} \mathrm{op}_t(b^-)(\gamma_1(w^-)) = -\mathrm{op}_t(b^-)(\gamma_1(w^+)) + \mathrm{op}_t(b^-)(k\gamma_0(w^+)) + \mathrm{op}_t(b^-)(\widetilde{G}_1^-) \\ \gamma_0(w^-) = \gamma_0(w^+) - \theta_{\varphi,\chi^0} \end{cases} \tag{3.53}$$

where $k = -i((\partial_{x_n}\varphi^-)_{|x_n=0} + (\partial_{x_n}\varphi^+)_{|x_n=0})$ and $\widetilde{G}_1^- = i(\partial_{x_n}\varphi^-)_{|x_n=0}\theta_{\varphi,\chi^0} + G_1$ that satisfies from estimate (3.48) that

$$|\widetilde{G}_1^-|_0 \leq C(h|v_{|x_n=0}|_0 + |\theta_\varphi|_0 + |\tilde{\theta}_\varphi|_0). \tag{3.54}$$

Then with combining (3.50) and (3.53) we obtain

$$\begin{aligned} \overbrace{(\mathrm{op}_t(1-a^-) - \mathrm{op}_t(b^-)\circ k)}^{=\mathrm{op}_t(\kappa)}(\gamma_0(w^+)) &= -\mathrm{op}_t(b^-)(\gamma_1(w^+)) \\ &\quad + \mathrm{op}_t(b^-)(\widetilde{G}_1^-) + \mathrm{op}_t(1-a^-)(\theta_{\varphi,\chi^0}) + G_2^-. \end{aligned} \tag{3.55}$$

Let $\chi_2(x,\xi') \in S_t^0$, satisfy the same requirement as χ^0 such taht

$$\begin{cases} \chi_2(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi^0) \\ \chi_1(x,\xi') \equiv 1 & \text{in neighborhood of } \mathrm{supp}(\chi_2). \end{cases}$$

Then from (3.51) the principal symbol of κ is given by

$$\kappa_{0\,|\mathrm{supp}(\chi_2)} = \frac{\rho^{-,+} - k}{\rho^{-,+} - \rho^{-,-}} \in S_t^0.$$

In $\mathrm{supp}(\chi_2)$ we have $\mathrm{Im}(\rho^{-,+}) \geq C$, $\mathrm{Im}(\rho^{-,-}) \leq -C$ and $\mathrm{Im}(-k) > 0$ by assumption (3.6), then κ is elliptic in $\mathrm{supp}(\chi_2)$. Hence, there exists $l \in S_t^0$, with $l = \sum_{j=0}^{M} h^j l_j$, $l_j \in S_t^{-j}$ and $l_0 = \frac{\chi_2}{\kappa_0}$, such that for M large we have

$$\mathrm{op}_t(l) \circ \mathrm{op}_t(\kappa) = \mathrm{op}_t(\chi_2) + h^{M+1}R_M;\ R_M \in \Psi_t^{-M-1}.$$

Thus from (3.55) we find

$$\gamma_0(w^+) = -\mathrm{op}_t(l) \circ \mathrm{op}_t(b^-)(\gamma_1(w^+)) + G_3 \tag{3.56}$$

with

$$G_3 = \mathrm{op}_t(l)\circ\mathrm{op}_t(b^-)\widetilde{G}_1^- + \mathrm{op}_t(l)\circ\mathrm{op}_t(1-a^-)\theta_{\varphi,\chi^0} + \mathrm{op}_t(l)G_2^- + \mathrm{op}_t(1-\chi_2)\gamma_0(w^+) - h^{M+1}R_M\gamma_0(w^+).$$

Since $\mathrm{supp}(1-\chi_2) \cap \mathrm{supp}(\chi^0) = \emptyset$, we obtain by estimates (3.52), (3.54) and Remark A.1 that

$$|G_3|_1 \leq C(h^{-\frac{1}{2}}\|P_\varphi^- v^-\|_0 + h^{\frac{1}{2}}\|v^-\|_1 + h^{\frac{3}{2}}|(D_{x_n}v^-)_{|x_n=0}|_0 + |\theta_{\varphi,\chi^0}|_1 + |\tilde{\theta}_\varphi|_0). \tag{3.57}$$

Thus by (3.56) we obtain

$$|\gamma_0(w^+)|_1 \leq C(|\gamma_1(w^+)|_0 + h^{-\frac{1}{2}}\|P_\varphi^- v^-\|_0 + h^{\frac{1}{2}}\|v\|_1 + h^{\frac{3}{2}}|(D_{x_n}v^-)_{|x_n=0}|_0 + |\theta_{\varphi,\chi^0}|_1 + |\tilde{\theta}_\varphi|_0). \tag{3.58}$$

Classical Carleman techniques for P_φ^+: As in proof of Proposition 3.3 and with keeping the same notations we get

$$h\|w^+\|_1^2 + \mathrm{Re}\left(h\mathcal{B}^+(w^+) + h^2\left((D_{x_n}w^+ + L_1^+w^+)_{|x_n=0}, L_0^+w^+_{|x_n=0}\right)_0\right) \leq C\|P_\varphi^+ w^+\|_0^2 \tag{3.59}$$

for h sufficiently small and

$$h^2 \left| \left((D_{x_n} w^+ + L_1^+ w^+)_{|x_n=0}, L_0^+ w^+_{|x_n=0} \right)_0 \right| \leq Ch^2 (|\gamma_1(w^+)|_0^2 + |\gamma_0(w^+)|_1^2). \tag{3.60}$$

From (3.35) and (3.56) we write

$$\mathcal{B}^+(w^+) = \big(M^* B^+ M \gamma_1(w^+), \gamma_1(w^+) \big)_0 + U(\gamma_1(w^+), G_3) \tag{3.61}$$

where

$$\left\{ \begin{array}{l} M = \left(\begin{array}{c} 1 \\ -\Lambda_t^1 \circ \mathrm{op}_t(l) \circ \mathrm{op}_t(b^-) \end{array} \right) \in \Psi_t^0 \\ U(\psi_1, \psi_2) = \left(B^+ M \psi_1, \left(\begin{array}{c} 0 \\ \Lambda_t^1 \psi_2 \end{array} \right) \right)_0 + \left(B^+ \left(\begin{array}{c} 0 \\ \Lambda_t^1 \psi_2 \end{array} \right), M\psi_1 \right)_0 \\ \qquad + \left(B^+ \left(\begin{array}{c} 0 \\ \Lambda_t^1 \psi_2 \end{array} \right), \left(\begin{array}{c} 0 \\ \Lambda_t^1 \psi_2 \end{array} \right) \right)_0 . \end{array} \right. \tag{3.62}$$

That satisfies from estimate (3.62) and Young inequality

(3.63)
$$|U(\gamma_1(w^+), G_3)| \leq \epsilon' |\gamma_1(w^+)|_0^2 + C_{\epsilon'} (h^{-1} \|P_\varphi^- v^-\|_0^2 + h\|v^-\|_1^2 + h^3 |(D_{x_n} v^-)_{|x_n=0}|_0^2 + |\theta_{\varphi,\chi^0}|_1^2 + |\tilde{\theta}_\varphi|_0^2).$$

From (3.37) and (3.62) we get

$$\sigma(M^* B^+ M) = \big(2(\partial_{x_n} \varphi^+) - 4 q_1^+ \mathrm{Re}(l_0 b_{-1}^-) - 2|l_0 b_{-1}^-|^2 (\partial_{x_n} \varphi^+) q_2^+ \big)_{|x_n=0} .$$

In $\mathrm{supp}(\chi^0)$ we have the following equalities

$$\left\{ \begin{array}{l} |l_0 b_{-1}^-|^{-2}_{|x_n=0} = |h\rho^{-,+}_{|x_n=0} - k|^2 = \Big((\mathrm{Re}(\rho^{-,+}))^2 + (\mathrm{Im}(\rho^{-,+}) + (\partial_{x_n} \varphi^-) + (\partial_{x_n} \varphi^+))^2 \Big)_{|x_n=0} \\ (\mathrm{Re}(l_0 b_{-1}^-) |l_0 b_{-1}^-|^{-2})_{|x_n=0} = \mathrm{Re}(\rho^{-,+})_{|x_n=0}. \end{array} \right.$$

Then Lemma 3.2 and assumptions (3.6) yields

$$\begin{aligned} &\sigma(M^* B^+ M) = 2|l_0 b_{-1}^-|^2 (\partial_{x_n} \varphi^+)_{|x_n=0} \big(|l_0 b_{-1}^-|^{-2} - 2 q_1^+ (\partial_{x_n} \varphi^+)^{-1} \mathrm{Re}(\rho^{-,+}) - q_2^+ \big)_{|x_n=0} = \\ &2|l_0 b_{-1}^-|^2 (\partial_{x_n} \varphi^+)_{|x_n=0} \Big(|l_0 b_{-1}^-|^{-2} - (\mathrm{Re}(\rho^{-,+}))^2 - \mu^+ + \big(q_1^+ (\partial_{x_n} \varphi^+)^{-1} - \mathrm{Re}(\rho^{-,+}) \big)^2 \Big)_{|x_n=0} \\ &\geq 2|l_0 b_{-1}^-|^2 (\partial_{x_n} \varphi^+)_{|x_n=0} \Big(\big((\partial_{x_n} \varphi^-) + (\partial_{x_n} \varphi^+) \big)^2 - \mu^+ \Big)_{|x_n=0} \geq C. \end{aligned}$$

Hence, Gårding inequality (Lemma A.2) yields for h sufficiently small

$$\mathrm{Re}(M^* B^+ M \gamma_1(w^+), \gamma_1(w^+))_0 \geq C |\gamma_1(w^+)|_0^2. \tag{3.64}$$

Now from (3.46), (3.58), (3.59), (3.60), (3.61), (3.63) and (3.64) we obtain for ϵ' and h sufficiently small

$$\begin{aligned} &h\|w^+\|_1^2 + h|\gamma_1(w^+)|_0^2 + h|\gamma_0(w^+)|_1^2 \leq \\ &\qquad C(\|P_\varphi v\|_0^2 + h^2 \|v\|_1^2 + h^4 |(D_{x_n} v^-)_{|x_n=0}|_0^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2). \end{aligned} \tag{3.65}$$

Note that as the proof of Proposition 3.2 we have by transmission conditions

$$|\gamma_0(w^-)|_1^2 + |\gamma_1(w^-)|_0^2 \leq C(|\gamma_0(w^+)|_1^2 + |\gamma_1(w^+)|_0^2 + h\|v\|_1^2 + |\theta_\varphi|_1^2 + |\tilde{\theta}_\varphi|_0^2). \tag{3.66}$$

Then from (3.65) and (3.66) we have

$$\begin{aligned} &h\|w^+\|_1^2 + h|\gamma_1(w^-)|_0^2 + h|\gamma_0(w^-)|_1^2 \leq \\ &\qquad C(\|P_\varphi v\|_0^2 + h^2 \|v\|_1^2 + h^4 |(D_{x_n} v^-)_{|x_n=0}|_0^2 + h|\theta_\varphi|_1^2 + h|\tilde{\theta}_\varphi|_0^2). \end{aligned} \tag{3.67}$$

We recall that

$$D_{x_n} \mathrm{op}_t(\chi^0) v = \mathrm{op}_t(\chi^0)(D_{x_n} v) + \underbrace{[D_{x_n}, \mathrm{op}_t(\chi^0)]}_{\in h\Psi_t^0} v.$$

Then the result will be achieved by combining the estimates (3.49), (3.65) and (3.67) as in proof of Proposition 3.2. ■

3.6 Boundary Carleman estimate

Let v_1, v_2, f_1 and f_2 satisfying the following boundary equations

$$\left\{\begin{array}{ll} B_1(x,D)(v_1) = f_1 & \text{in } \Omega_1 \\ B_2(x,D)(v_2) = f_2 & \text{in } \Omega_1 \\ hv_1 = v_2 + \theta & \text{on } S \\ h\partial_\nu v_1 + \partial_\nu v_2 = \tilde{\theta} & \text{on } S \end{array}\right. \tag{3.68}$$

where $B_1(x,D)$ and $B_2(x,D)$ defined a differential operators with C^∞-coefficients gived by

$$\left\{\begin{array}{l} B_1(x,D)(v_1) = -\Delta v_1 - \dfrac{1}{h} v_1 \\ \\ B_2(x,D)(v_2) = -\Delta v_2 + \dfrac{1}{h} v_2 \end{array}\right.$$

with h is a small pseudo-differential parameter. This mean that the trace of v_1 and v_2 at the boundary S, as well as the their the normal flux, with some coefficients terms, are respectively equal and opposed modulo some error terms θ and $\tilde{\theta}$ respectively. The aim of this section is to find a local Carleman estimate for a such problem.

In a neighborhood V of a point $z = (z', z_n) \in S$, we denote by x_n the variable that is normal to the interface and by x' the remainding variables, i.e $x = (x', x_n)$ in particular $z = (z', 0)$. The edge is given now by $S = \{x_n = 0\}$. In a sufficient small neighborhood $V \subset \mathbb{R}^n$ of z we place our selves in normal geodesic coordinates. For convience, we shall take the neighborhood V of the form $V = V_{z'} \times]0, \epsilon[$ where $V_{z'}$ is a sufficient small neighborhood of z'. In such coordinate system the operators B_1 and B_2 take the following forms

$$\left\{\begin{array}{l} B_1(x,D) := -\partial_{x_n}^2 + R_1(x, \partial_{x'}/i) - \dfrac{1}{h} \\ \\ B_2(x,D) := -\partial_{x_n}^2 + R_2(x, \partial_{x'}/i) + \dfrac{1}{h} \end{array}\right.$$

where $R_1(x,\xi')$ and $R_2(x,\xi')$ are a second ordre polynomials in ξ' with coefficients in $\mathbb{R}$ with principal symbol respectively $r_1(x,\xi')$ and $r_2(x,\xi')$ that satisfy

$$\left\{\begin{array}{ll} r_1(x,\xi') \geq C|\xi'|^2 & \forall\, (x', x_n) \in V_{z'} \times]0,\epsilon[,\ \forall\, \xi' \in \mathbb{R}^{n-1} \\ r_2(x,\xi') \geq C|\xi'|^2 & \forall\, (x', x_n) \in V_{z'} \times]0,\epsilon[,\ \forall\, \xi' \in \mathbb{R}^{n-1}. \end{array}\right.$$

Let ψ_1 and ψ_2 two weight functions and we define the two adjoint operators

$$B_{\psi_1} = h^2 \mathrm{e}^{\psi_1/h} B_1 \mathrm{e}^{-\psi_1/h} \quad \text{and} \quad B_{\psi_2} = h^2 \mathrm{e}^{\psi_2/h} B_2 \mathrm{e}^{-\psi_2/h}$$

with principal symbols respectively

$$\left\{\begin{array}{l} b_{\psi_1}(x,\xi) = (\xi_n + i(\partial_{x_n}\psi_1))^2 + r_1(x, \xi' + i(\partial_{x'}\psi_1)) \\ b_{\psi_2}(x,\xi) = (\xi_n + i(\partial_{x_n}\psi_2))^2 + r_2(x, \xi' + i(\partial_{x'}\psi_2)). \end{array}\right.$$

Assumption 3.2 *The weight functions ψ_1 and ψ_2 are in $\mathscr{C}^\infty(\overline{V})$ and satisfies*

1. $|\nabla\psi_1|(x) > 0$ *and* $|\nabla\psi_2|(x) > 0$ *in* $\overline{V}$.
2. $\psi_1(x', 0) = \psi_2(x', 0)$.
3. $$\left\{\begin{array}{l} (\partial_{x_n}\psi_1)(x',0) > 0 \\ (\partial_{x_n}\psi_1)(x',0) + (\partial_{x_n}\psi_2)(x',0) > 0 \qquad \textit{for all } x' \in V_{z'}. \\ (\partial_{x_n}\psi_1)^2(x',0) - (\partial_{x_n}\psi_2)^2(x',0) > 0 \end{array}\right.$$

4. The sub-ellipticity condition:

$$\left\{\begin{array}{l}\forall\,(x,\xi)\in\overline{V}\times\mathbb{R}^n;\ b_{\psi_1}(x,\xi)=0\Longrightarrow\{\mathrm{Re}(b_{\psi_1}),\mathrm{Im}(b_{\psi_1})\}(x,\xi)>0\\ \forall\,(x,\xi)\in\overline{V}\times\mathbb{R}^n;\ b_{\psi_2}(x,\xi)=0\Longrightarrow\{\mathrm{Re}(b_{\psi_2}),\mathrm{Im}(b_{\psi_2})\}(x,\xi)>0.\end{array}\right.$$

From the assumptions made on the weight functions ψ_1 and ψ_2 we shall obtain the following local Carleman estimate as we done in the previous sections.

Theorem 3.2 *Let K be a compact subset of V and ψ_1 and ψ_2 a weight functions satisfying Assumption 3.2, then there exist $C>0$ and $h_0>0$ such that*

$$\begin{aligned}&h^3\|\mathrm{e}^{\psi_1/h}v_1\|_0^2+h^5\|\mathrm{e}^{\psi_1/h}\nabla v_1\|_0^2+h^3|\mathrm{e}^{\psi_1/h}v_{1\,|x_n=0}|_0^2+h^5|\mathrm{e}^{\psi_1/h}\nabla_{x'}v_{1\,|x_n=0}|_0^2\\ &+h^5|\mathrm{e}^{\psi_1/h}\partial_{x_n}v_{1\,|x_n=0}|_0^2+h\|\mathrm{e}^{\psi_2/h}v_2\|_0^2+h^3\|\mathrm{e}^{\psi_2/h}\nabla v_2\|_0^2+h|\mathrm{e}^{\psi_2/h}v_{2\,|x_n=0}|_0^2\\ &+h^3|\mathrm{e}^{\psi_2/h}\nabla_{x'}v_{2\,|x_n=0}|_0^2+h^3|\mathrm{e}^{\psi_2/h}\partial_{x_n}v_{2\,|x_n=0}|_0^2\le C(h^6\|\mathrm{e}^{\psi_1/h}f_1\|_0^2+h^4\|\mathrm{e}^{\psi_2/h}f_2\|_0^2\\ &+h|\mathrm{e}^{\psi_1/h}\theta|_0^2+h^3|\mathrm{e}^{\psi_1/h}\nabla_{x'}\theta|_0^2+h^3|\mathrm{e}^{\psi_1/h}\tilde{\theta}|_0^2)\end{aligned}$$

for all $0<h\le h_0$, v_1, v_2, f_1 and f_2 satisfying (3.68) where $v_{1\,|\overline{\mathbb{R}^n_+}}$, $v_{2\,|\overline{\mathbb{R}^n_+}}\in\mathscr{C}_c^\infty(K)$.

4 Global Carleman estimates and construction of the weight functions

The purpose of this section is to state two global Carleman estimates, the first is gived for two second orde operators in Ω_1 and Ω_2 respectively with transmission conditions on the interface S, and the second one is gived for two second order and elliptic operators both of them are in Ω_2 with functions for each operator which are linked on the interface S by a boundary condions. As a second step we will try to construct a very special weight functions for each of the two Carleman estimates cited before so as to able to combining the two estimates in the next section.

We consider the following value boundary transmission problem

$$(4.1)\qquad\left\{\begin{array}{lll}-\Delta u_1=f_1 & \text{in} & \Omega_1\\ -\Delta u_2-\dfrac{1}{h^2}u_2=f_2 & \text{in} & \Omega_2\\ u_1=u_2+\theta & \text{on} & S\\ \partial_\nu u_1=\partial_\nu u_2+\tilde{\theta} & \text{on} & S\\ u_2=0 & \text{on} & \Gamma_2.\end{array}\right.$$

We define the operator

$$A(x,D)=\left\{\begin{array}{ll}A_1(x,D)=-\Delta & \text{in }\Omega_1\\ A_2(x,D)=-\Delta-\dfrac{1}{h^2} & \text{in }\Omega_2,\end{array}\right.$$

and the adjoint operator

$$A_\varphi(x,D)=h^2\mathrm{e}^{\varphi/h}A\mathrm{e}^{-\varphi/h},$$

with principal symbol

$$a_\varphi(x,\xi)=\left\{\begin{array}{ll}|\xi+i\nabla\varphi_1|^2 & \text{in }\Omega_1\\ |\xi+i\nabla\varphi_2|^2-1 & \text{in }\Omega_2\end{array}\right.$$

where $\varphi=(\varphi_1,\varphi_2)$ is a weight function.

Assumption 4.1 *Let $\varphi=(\varphi_1,\varphi_2)$ be a continuous weight function in $\overline{\Omega}$ such that*

1. $\varphi_1\in\mathscr{C}^\infty(\overline{\Omega}_1)$ *and* $\varphi_2\in\mathscr{C}^\infty(\overline{\Omega}_2)$
2. $|\nabla\varphi|(x)>0$ *in* $\overline{\Omega}$

3. $\begin{cases} \partial_\nu \varphi_{1\,|S} < 0 \\ \partial_\nu \varphi_{1\,|S} - \partial_\nu \varphi_{2\,|S} < 0 \\ (\partial_\nu \varphi_{1\,|S})^2 - (\partial_\nu \varphi_{2\,|S})^2 > 1 \end{cases}$

4. $\partial_\nu \varphi_{1\,|\Gamma_1} \neq 0$ *and* $\partial_\nu \varphi_{2\,|\Gamma_2} < 0$

5. The sub-ellipticity condition:

$$\forall\, (x,\xi) \in \overline{\Omega} \times \mathbb{R}^n;\ a_\varphi(x,\xi) = 0 \Longrightarrow \{\mathrm{Re}(a_\varphi), \mathrm{Im}(a_\varphi)\}(x,\xi) > 0. \tag{4.2}$$

From Theorem 3.1, [LR97, Proposition 1] and [LR95, Proposition 2] we get the following Carleman estimate.

Theorem 4.1 *Let* $\varphi = (\varphi_1, \varphi_2)$ *be a weight function that satisfy Assumption 4.1, then there exist* $C > 0$ *and* $h_0 > 0$ *such that*

$$\begin{aligned} &h\|\mathrm{e}^{\varphi_1/h}u_1\|^2_{L^2(\Omega_1)} + h^3\|\mathrm{e}^{\varphi_1/h}\nabla u_1\|^2_{L^2(\Omega_1)} + h|\mathrm{e}^{\varphi_1/h}u_1|^2_{L^2(S)} + h^3|\mathrm{e}^{\varphi_1/h}\nabla u_1|^2_{L^2(S)} \\ &+h\|\mathrm{e}^{\varphi_2/h}u_2\|^2_{L^2(\Omega_2)} + h^3\|\mathrm{e}^{\varphi_2/h}\nabla u_2\|^2_{L^2(\Omega_2)} + h|\mathrm{e}^{\varphi_2/h}u_2|^2_{L^2(S)} + h^3|\mathrm{e}^{\varphi_2/h}\nabla u_2|^2_{L^2(S)} \\ &\le C(h^4\|\mathrm{e}^{\varphi_1/h}f_1\|^2_{L^2(\Omega_1)} + h^4\|\mathrm{e}^{\varphi_2/h}f_2\|^2_{L^2(\Omega_2)} + h|\mathrm{e}^{\varphi/h}\theta|^2_{L^2(S)} + h^3|\mathrm{e}^{\varphi/h}\nabla\theta|^2_{L^2(S)} \\ &\qquad +h^3|\mathrm{e}^{\varphi/h}\tilde{\theta}|^2_{L^2(S)} + h|\mathrm{e}^{\varphi_1/h}u_1|^2_{L^2(\Gamma_1)} + h^3|\mathrm{e}^{\varphi_1/h}\partial_\nu u_1|^2_{L^2(\Gamma_1)}) \end{aligned}$$

for all $0 < h \le h_0$, $u = (u_1, u_2) \in \mathscr{C}^\infty(\overline{\Omega}_1) \times \mathscr{C}^\infty(\overline{\Omega}_2)$ *and* $f = (f_1, f_2)$ *satisfying to the system* (4.1).

We consider now the following boundary value problem

$$\begin{cases} -\Delta v_1 - \dfrac{1}{h} v_1 = g_1 & \text{in} \quad \Omega_1 \\ \\ -\Delta v_2 + \dfrac{1}{h} v_2 = g_2 & \text{in} \quad \Omega_1 \\ h v_1 = v_2 + \theta & \text{on} \quad S \\ h\partial_\nu v_1 + \partial_\nu v_2 = \tilde{\theta} & \text{on} \quad S. \end{cases} \tag{4.3}$$

We define in Ω_2 the operators

$$\begin{cases} B_1(x,D) = -\Delta - \dfrac{1}{h} \\ \\ B_2(x,D) = -\Delta + \dfrac{1}{h}, \end{cases}$$

and the adjoint operators

$$B_{\psi_1}(x,D) = h^2 \mathrm{e}^{\psi_1/h} B_1 \mathrm{e}^{-\psi_1/h} \quad \text{and} \quad B_{\psi_2}(x,D) = h^2 \mathrm{e}^{\psi_2/h} B_2 \mathrm{e}^{-\psi_2/h}$$

with principal symbol respectively

$$b_{\psi_1}(x,\xi) = |\xi + i\nabla\psi_1|^2 \quad \text{and} \quad b_{\psi_2}(x,\xi) = |\xi + i\nabla\psi_2|^2$$

where ψ_1, ψ_2 are two weight functions.

Assumption 4.2 *Let* ψ_1 *and* ψ_2 *be two continuous weight function in* $\overline{\Omega_1}$ *such that*

1. $\psi_1 \in \mathscr{C}^\infty(\overline{\Omega}_1)$ *and* $\psi_2 \in \mathscr{C}^\infty(\overline{\Omega}_1)$ *and* $\psi_{1\,|S} = \psi_{2\,|S}$
2. $|\nabla\psi_1|(x) > 0$ *in* $\overline{\Omega}_1$ *and* $|\nabla\psi_2|(x) > 0$ *in* $\overline{\Omega}_1$
3. $\begin{cases} \partial_\nu \psi_{1\,|S} < 0 \\ \partial_\nu \psi_{1\,|S} + \partial_\nu \psi_{2\,|S} < 0 \\ (\partial_\nu \psi_{1\,|S})^2 - (\partial_\nu \psi_{2\,|S})^2 > 0 \end{cases}$
4. $\partial_\nu \psi_{1\,|\Gamma_1} \neq 0$ *and* $\partial_\nu \psi_{2\,|\Gamma_1} \neq 0$

5. *The sub-ellipticity condition:*

$$\forall\,(x,\xi)\in\overline{\Omega}_1\times\mathbb{R}^n;\ b_{\psi_1}(x,\xi)=0\Longrightarrow\{\mathrm{Re}(b_{\psi_1}),\mathrm{Im}(b_{\psi_1})\}(x,\xi)>0 \tag{4.4}$$

$$\forall\,(x,\xi)\in\overline{\Omega}_1\times\mathbb{R}^n;\ b_{\psi_2}(x,\xi)=0\Longrightarrow\{\mathrm{Re}(b_{\psi_2}),\mathrm{Im}(b_{\psi_2})\}(x,\xi)>0. \tag{4.5}$$

From Theorem 3.2 and [LR97, Proposition 1] we get the following Carleman estimate.

Theorem 4.2 *Let ψ_1 and ψ_2 be two weight functions that satisfies Assumption 4.2, then there exist $C>0$ and $h_0>0$ such that*

$$\begin{aligned}
&h^3\|\mathrm{e}^{\psi_1/h}v_1\|^2_{L^2(\Omega_1)}+h^5\|\mathrm{e}^{\psi_1/h}\nabla v_1\|^2_{L^2(\Omega_1)}+h^3|\mathrm{e}^{\psi_1/h}v_1|^2_{L^2(S)}+h^5|\mathrm{e}^{\psi_1/h}\nabla v_1|^2_{L^2(S)}\\
&+\ h\|\mathrm{e}^{\psi_2/h}v_2\|^2_{L^2(\Omega_1)}+h^3\|\mathrm{e}^{\psi_2/h}\nabla v_2\|^2_{L^2(\Omega_1)}+h|\mathrm{e}^{\psi_2/h}v_2|^2_{L^2(S)}+h^3|\mathrm{e}^{\psi_2/h}\nabla v_2|^2_{L^2(S)}\leq\\
&C(h^6\|\mathrm{e}^{\psi_1/h}g_1\|^2_{L^2(\Omega_1)}+h^4\|\mathrm{e}^{\psi_2/h}g_2\|^2_{L^2(\Omega_1)}+h|\mathrm{e}^{\varphi/h}\theta|^2_{L^2(S)}+h^3|\mathrm{e}^{\varphi/h}\nabla\theta|^2_{L^2(S)}+h^3|\mathrm{e}^{\varphi/h}\tilde{\theta}|^2_{L^2(S)}\\
&+\ h^3|\mathrm{e}^{\psi_1/h}v_1|^2_{L^2(\Gamma_1)}+h^5|\mathrm{e}^{\psi_1/h}\partial_\nu v_1|^2_{L^2(\Gamma_1)}+h|\mathrm{e}^{\psi_2/h}v_2|^2_{L^2(\Gamma_1)}+h^3|\mathrm{e}^{\psi_2/h}\partial_\nu v_2|^2_{L^2(\Gamma_1)})
\end{aligned}$$

for all $0<h\leq h_0$, v_1 and v_2 are in $\mathscr{C}^\infty(\overline{\Omega}_1)$ and g_1 and g_2 that satisfies the system (4.3).

For the first Carleman estimate now we will try to find two phases $\varphi_1=(\varphi_{1,1},\varphi_{1,2})$ and $\varphi_2=(\varphi_{2,1},\varphi_{2,2})$ defined in $\overline{\Omega}$ satisfies the Hörmander's condition except in a finite number of balls where one of them do not satisfy this condition the second does and is strictly greater. For the second Carleman estimate we will try to find four phases $\psi_{1,1}$, $\psi_{1,2}$, $\psi_{2,1}$ and $\psi_{2,2}$ defined in $\overline{\Omega}_1$ satisfying the Hörmander's condition exept in a finte number of balls where $\psi_{1,1}$ or $\psi_{2,1}$ (resp. $\psi_{1,2}$ or $\psi_{2,2}$) do not satisfy this condition the other does and is strictly greater.

From [Has13, Proposition 3.2] and [Bur98, Proposition 3.2] we can find two $\mathscr{C}^\infty$ functions $\varphi_{1,1}$ and $\varphi_{2,1}$ in $\overline{\Omega}_1$ (resp. $\varphi_{1,2}$ and $\varphi_{2,2}$ in $\overline{\Omega}_2$) such that for $i,k\in\{1,2\}$ there exists a finite number of points $x^k_{1,j_1}\in\Omega_k$ and $x^k_{2,j_2}\in\Omega_k$ for $j_i=1,\dots,N_{i,k}$ and $\epsilon>0$ such that $B(x^k_{i,j_i},2\epsilon)\subset\Omega_k$ and $B(x^k_{1,j_1},2\epsilon)\bigcap B(x^k_{2,j_2},2\epsilon)=\emptyset$ and in $B(x^k_{i,j_i},2\epsilon)$ we have $\varphi_{\sigma(i),k}>\varphi_{i,k}$ where σ is the permutation of the set $\{1,2\}$ different from the identity, where we denote by $U_{i,k}=\Omega_k\bigcap\left(\bigcup_{j_i=1}^{N_{i,k}}B(x^k_{i,j_i},2\epsilon)\right)^c$, verifying that $|\nabla\varphi_{1,1}|>0$ in $U_{1,1}$, $|\nabla\varphi_{2,1}|>0$ in $U_{2,1}$, $\partial_\nu\varphi_{i,1\,|\Gamma_1}\neq0$, $\partial_\nu\varphi_{i,1\,|S}\neq0$, $\partial_\nu\varphi_{i,2\,|\Gamma_2}\neq0$, $\partial_\nu\varphi_{i,2\,|S}\neq0$ and φ_1 and φ_2 verifying the Hörmander's condition (4.2) in $U_{1,1}\cup U_{1,2}$ and $U_{2,1}\cup U_{2,2}$ respectively. By construction we can suppose that $\varphi_{i,1\,|S}=\varphi_{i,2\,|S}$, $\partial_\nu\varphi_{i,1\,|S}-\partial_\nu\varphi_{i,2\,|S}<0$ and $(\partial_\nu\varphi_{i,1\,|S})^2-(\partial_\nu\varphi_{i,2\,|S})^2>0$.

For the same arguments as above we can find two $\mathscr{C}^\infty$ functions $\psi_{1,1}$ and $\psi_{2,1}$ (resp. $\psi_{1,2}$ and $\psi_{2,2}$) in $\overline{\Omega}_1$ such that for $i,k\in\{1,2\}$ there exists a finite number of points $y^k_{i,j_i}\in\Omega_1$ for $j_i=1,\dots,M_{i,k}$ and $\epsilon>0$ such that $B(y^k_{i,j_i},2\epsilon)\subset\Omega_1$ and $B(y^k_{1,j_1},2\epsilon)\bigcap B(y^k_{2,j_2},2\epsilon)=\emptyset$ and in $B(y^k_{i,j_i},2\epsilon)$ we have $\psi_{\sigma(i),k}>\psi_{i,k}$, where we denote by $V_{i,k}=\Omega_1\bigcap\left(\bigcup_{j_i=1}^{M_{i,k}}B(y^k_{i,j_i},2\epsilon)\right)^c$, verifying that $|\nabla\psi_{1,1}|>0$ in $\overline{V}_{1,1}$, $|\nabla\psi_{2,1}|>0$ in $\overline{V}_{2,1}$, $\partial_\nu\psi_{i,k\,|\Gamma_1}\neq0$, $\partial_\nu\psi_{i,1\,|S}<0$ and $\psi_{1,1}$ and $\psi_{2,1}$ verifying the Hörmander's condition (4.2) in $V_{1,1}$ and $V_{1,2}$ respectively and $\psi_{1,2}$ and $\psi_{2,2}$ verifying the Hörmander's condition (4.4) in $V_{2,1}$ and $V_{2,2}$ respectively. And we can suppose that $\psi_{i,1\,|S}=\psi_{i,2\,|S}$, $\partial_\nu\psi_{i,1\,|S}+\partial_\nu\psi_{i,2\,|S}<0$ and $(\partial_\nu\psi_{i,1\,|S})^2-(\partial_\nu\psi_{i,2\,|S})^2>0$.

Finally, to link the two construction we can suppose that $\varphi_{1,1}=\psi_{1,2}$ and $\varphi_{2,1}=\psi_{2,2}$. This assumption will play a very important role in the proof of Theorem 1.2 in the next section and in particular the trace of $\varphi_{1,1}$ and $\psi_{1,2}$ (resp. $\varphi_{2,1}$ and $\psi_{2,2}$) on the edge of Ω_1 are equal.

5 Resolvent Estimate

The aim of this section is to prove the resolvent estimate, precisely we prove for some constante $C>0$ that

$$\|(\mathcal{A}-i\mu\,\mathrm{Id})^{-1}\|_{\mathcal{L}(\mathcal{H})}\leq C\mathrm{e}^{C|\mu|},$$

for $|\mu|$ large enough.

The main idea consist to derived from the plate-wave equation two systems of second order of the same form as (4.1) and (4.3) respectively and to apply to them the appropriate Carleman estimate established in the previous section. And with the choice of the weight functions constructed in section 4 the result follow from the combination of these two last estimates.

Let $(F, G) \in \mathcal{H}$ with $F = (f_1, f_2)$ and $G = (g_1, g_2)$ and $(u, v) \in \mathcal{D}(A)$ with $u = (u_1, u_2)$ and $v = (v_1, v_2)$ such that

$$(\mathcal{A} - i\mu) \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} F \\ G \end{pmatrix},$$

which can be written as follows

$$\left\{ \begin{array}{lll} v_1 - i\mu u_1 = f_1 & \text{in} & \Omega_1 \\ v_2 - i\mu u_2 = f_2 & \text{in} & \Omega_2 \\ -\Delta^2 u_1 - i\mu v_1 = g_1 & \text{in} & \Omega_1 \\ \Delta u_2 - i\mu v_2 = g_2 & \text{in} & \Omega_2 \\ u_1 + u_2 = 0 & \text{on} & S \\ \partial_\nu \Delta u_1 = \partial_\nu u_2 & \text{on} & S \\ \Delta u_1 = 0 & \text{on} & S \\ \Delta u_1 = -a\partial_\nu v_1 & \text{on} & \Gamma_1 \\ \partial_\nu \Delta u_1 = b v_1 & \text{on} & \Gamma_1 \\ u_2 = 0 & \text{on} & \Gamma_2. \end{array} \right. \tag{5.1}$$

We set now the following functions

$$\Phi = \left\{ \begin{array}{ll} \Phi_1 = -g_1 - i\mu f_1 & \text{in } \Omega_1 \\ \Phi_2 = -g_2 - i\mu f_2 & \text{in } \Omega_2 \end{array} \right. \quad \textit{and} \quad \left\{ \begin{array}{ll} \phi_1 = -a\partial_\nu f_1 & \text{on } \Gamma_1 \\ \phi_2 = b f_1 & \text{on } \Gamma_1. \end{array} \right. \tag{5.2}$$

The system (5.1) is written now as follows

$$\left\{ \begin{array}{lll} v_1 = f_1 + i\mu u_1 & \text{in} & \Omega_1 \\ v_2 = f_2 + i\mu u_2 & \text{in} & \Omega_2 \\ \Delta^2 u_1 - \mu^2 u_1 = \Phi_1 & \text{in} & \Omega_1 \\ \Delta u_2 + \mu^2 u_2 = -\Phi_2 & \text{in} & \Omega_2 \\ u_1 + u_2 = 0 & \text{on} & S \\ \partial_\nu \Delta u_1 = \partial_\nu u_2 & \text{on} & S \\ \Delta u_1 = 0 & \text{on} & S \\ \Delta u_1 + i\mu a \partial_\nu u_1 = \phi_1 & \text{on} & \Gamma_1 \\ \partial_\nu \Delta u_1 - i\mu b u_1 = \phi_2 & \text{on} & \Gamma_1 \\ u_2 = 0 & \text{on} & \Gamma_2. \end{array} \right. \tag{5.3}$$

Integrating by parts then we obtain

$$\begin{aligned} \langle \Phi, u \rangle &= \langle \Delta^2 u_1, u_1 \rangle_{L^2(\Omega_1)} - \langle \Delta u_2, u_2 \rangle_{L^2(\Omega_2)} - \mu^2 (\|u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L^2(\Omega_2)}) \\ &= \|\Delta u_1\|^2_{L^2(\Omega_1)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} - \mu^2 (\|u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L^2(\Omega_2)}) - \langle \phi_1, \partial_\nu u_1 \rangle_{L^2(\Gamma_1)} \\ &\quad + \langle \phi_2, u_1 \rangle_{L^2(\Gamma_1)} + i\mu (\langle a\partial_\nu u_1, \partial_\nu u_1 \rangle_{L^2(\Gamma_1)} + \langle b u_1, u_1 \rangle_{L^2(\Gamma_1)}). \end{aligned} \tag{5.4}$$

Keeping only the imaginary part of (5.4) then we get

$$\begin{aligned} |\mu| (\|a^{\frac{1}{2}} \partial_\nu u_1\|^2_{L^2(\Gamma_1)} + \|b^{\frac{1}{2}} u_1\|_{L^2(\Gamma_1)})^2 \leq C(&\|\Phi_1\|_{L^2(\Omega_1)} . \|u_1\|_{L^2(\Omega_1)} + \|\Phi_2\|_{L^2(\Omega_2)} . \|u_2\|_{L^2(\Omega_2)} \\ &+ |\phi_1|_{L^2(\Gamma_1)} . |\partial_\nu u_1|_{L^2(\Gamma_1)} + |\phi_2|_{L^2(\Gamma_1)} . |u_1|_{L^2(\Gamma_1)}). \end{aligned} \tag{5.5}$$

We denote now by

$$\left\{ \begin{array}{l} y_1 = \Delta u_1 \\ y_2 = u_2 \end{array} \right. \tag{5.6}$$

that are form system (5.3) solution of the following transmission boundary equation

$$\left\{\begin{array}{lll} \Delta y_1 = \Phi_1 + \mu^2 u_1 & \text{in} & \Omega_1 \\ \Delta y_2 + \mu^2 y_2 = -\Phi_2 & \text{in} & \Omega_2 \\ y_1 = y_2 + u_1 & \text{on} & S \\ \partial_\nu y_1 = \partial_\nu y_2 & \text{on} & S \\ y_1 = \phi_1 - i\mu a \partial_\nu u_1 & \text{on} & \Gamma_1 \\ \partial_\nu y_1 = \phi_2 + i\mu b u_1 & \text{on} & \Gamma_1 \\ y_2 = 0 & \text{on} & \Gamma_2. \end{array}\right. \tag{5.7}$$

We denote also by

$$\left\{\begin{array}{l} z_1 = \Delta u_1 - |\mu| u_1 \\ z_2 = u_1 \end{array}\right. \tag{5.8}$$

that are form system (5.3) solution of the following transmission boundary equation

$$\left\{\begin{array}{lll} \Delta z_1 + |\mu| z_1 = \Phi_1 & \text{in} & \Omega_1 \\ \Delta z_2 - |\mu| z_2 = z_1 & \text{in} & \Omega_2 \\ \dfrac{1}{|\mu|} z_1 = z_2 & \text{on} & S \\ & & \\ \dfrac{1}{|\mu|}\partial_\nu z_1 + \partial_\nu z_2 = \dfrac{1}{|\mu|}\partial_\nu u_2 & \text{on} & S \\ & & \\ z_1 = \phi_1 - i\mu a \partial_\nu u_1 - |\mu| u_1 & \text{on} & \Gamma_1 \\ \partial_\nu z_1 = \phi_2 + i\mu b u_1 - |\mu| \partial_\nu u_1 & \text{on} & \Gamma_1. \end{array}\right. \tag{5.9}$$

Noting that in the sequel we are using the notations of the previous section, and we denote henceforth $h = \dfrac{1}{|\mu|}$. We will now focus to the system (5.7). Let $\varphi_1 = (\varphi_{1,1}, \varphi_{1,2})$ and $\varphi_2 = (\varphi_{2,1}, \varphi_{2,2})$ satisfying the conclusion of the section 4. Let $\chi_1 = (\chi_{1,1}, \chi_{1,2})$ and $\chi_2 = (\chi_{2,1}), \chi_{2,2})$ two cutt-off functions equal to one in $\left(\bigcup_{\substack{k=1,2 \\ j_1=1}}^{N_{1,k}} B(x^k_{1,j_1}, 2\epsilon)\right)^c$ and $\left(\bigcup_{\substack{k=1,2 \\ j_2=1}}^{N_{2,k}} B(x^k_{1,j_2}, 2\epsilon)\right)^c$ respectively and supported respectively in $\left(\bigcup_{\substack{k=1,2 \\ j_1=1}}^{N_{1,k}} B(x^k_{1,j_1}, \epsilon)\right)^c$ and $\left(\bigcup_{\substack{k=1,2 \\ j_2=1}}^{N_{2,k}} B(x^k_{1,j_2}, \epsilon)\right)^c$ (in order to eliminate the critical points of the phases functions φ_1 and φ_2).

Then for $r = 1, 2$, where we are noting by $\tilde{y}_{r,1} = \chi_{r,1} y_1$ and $\tilde{y}_{r,2} = \chi_{r,1} y_2$, we obtain from the system (5.7) the following equation

$$\left\{\begin{array}{lll} \Delta \tilde{y}_{r,1} = \Psi_{r,1} & \text{in} & \Omega_1 \\ \Delta \tilde{y}_{r,2} + \dfrac{1}{h^2}\tilde{y}_2 = \Psi_{r,2} & \text{in} & \Omega_2 \\ \tilde{y}_{r,1} = \tilde{y}_{r,2} + u_1 & \text{on} & S \\ \partial_\nu \tilde{y}_{r,1} = \partial_\nu \tilde{y}_{r,2} & \text{on} & S \\ \tilde{y}_{r,1} = \phi_1 - i\mu a \partial_\nu u_1 & \text{on} & \Gamma_1 \\ \tilde{y}_{r,2} = \phi_2 + i\mu b u_1 & \text{on} & \Gamma_1 \\ \tilde{y}_{r,2} = 0 & \text{on} & \Gamma_2, \end{array}\right. \tag{5.10}$$

where

$$\left\{\begin{array}{ll} \Psi_{r,1} = [\Delta, \chi_{r,1}] y_1 + \chi_{r,1}.\Phi_1 + \dfrac{1}{h^2}\chi_{r,1}.u_1 & \text{in } \Omega_1 \\ \Psi_{r,2} = [\Delta, \chi_{r,2}] y_2 - \chi_{r,2}.\Phi_2 & \text{in } \Omega_2. \end{array}\right. \tag{5.11}$$

Applying now Carleman estimate of Theorem 4.1 to the system (5.10) then for $r = 1, 2$ we obtain

$$\begin{aligned} h\|\mathrm{e}^{\varphi_{r,1}/h}\tilde{y}_{r,1}\|^2_{L^2(U_{r,1})} + h\|\mathrm{e}^{\varphi_{r,2}/h}\tilde{y}_{r,2}\|^2_{L^2(U_{r,2})} + h^3\|\mathrm{e}^{\varphi_{r,1}/h}\nabla\tilde{y}_{r,1}\|^2_{L^2(U_{r,1})} \\ +h^3\|\mathrm{e}^{\varphi_{r,2}/h}\nabla\tilde{y}_{r,2}\|^2_{L^2(U_{r,2})} + h^3|\mathrm{e}^{\varphi_{r,2}/h}\partial_\nu\tilde{y}_{r,2}|^2_{L^2(S)} \leq C(h^4\|\mathrm{e}^{\varphi_{r,1}/h}\Psi_{r,1}\|^2_{L^2(U_{r,1})} \\ +h^4\|\mathrm{e}^{\varphi_{r,2}/h}\Psi_{r,2}\|^2_{L^2(U_{r,2})} + h|\mathrm{e}^{\varphi_{r,1}/h}u_1|^2_{L^2(S)} + h^3|\mathrm{e}^{\varphi_{r,1}/h}\nabla u_1|^2_{L^2(S)} + h|\mathrm{e}^{\varphi_{r,1}/h}\phi_1|^2_{L^2(\Gamma_1)} \\ +h^3|\mathrm{e}^{\varphi_{r,1}/h}\phi_2|^2_{L^2(\Gamma_1)} + h^{-1}|\mathrm{e}^{\varphi_{r,1}/h}a\partial_\nu u_1|^2_{L^2(\Gamma_1)} + h|\mathrm{e}^{\varphi_{r,1}/h}bu_1|^2_{L^2(\Gamma_1)}). \end{aligned}$$

Then by the expression of $\Psi_{r,1}$ and $\Psi_{r,2}$ in (5.11) we get

$$\begin{aligned} h\|\mathrm{e}^{\varphi_{r,1}/h}\tilde{y}_{r,1}\|^2_{L^2(U_{r,1})} + h\|\mathrm{e}^{\varphi_{r,2}/h}\tilde{y}_{r,2}\|^2_{L^2(U_{r,2})} + h^3\|\mathrm{e}^{\varphi_{r,1}/h}\nabla\tilde{y}_{r,1}\|^2_{L^2(U_{r,1})} \\ +h^3\|\mathrm{e}^{\varphi_{r,2}/h}\nabla\tilde{y}_{r,2}\|^2_{L^2(U_{r,2})} + h^3|\mathrm{e}^{\varphi_{r,2}/h}\partial_\nu\tilde{y}_{r,2}|^2_{L^2(S)} \leq C(h^4\|\mathrm{e}^{\varphi_{r,1}/h}\Phi_1\|^2_{L^2(U_{r,1})} \\ +h^4\|\mathrm{e}^{\varphi_{r,2}/h}\Phi_2\|^2_{L^2(U_{r,2})} + h^4\|\mathrm{e}^{\varphi_{r,1}/h}[\Delta, \chi_{r,1}]y_1\|^2_{L^2(U_{r,1})} + h^4\|\mathrm{e}^{\varphi_{r,2}/h}[\Delta, \chi_{r,2}]y_2\|^2_{L^2(U_{r,2})} \\ +\|\mathrm{e}^{\varphi_{r,1}/h}u_1\|^2_{L^2(U_{r,1})} + h|\mathrm{e}^{\varphi_{r,1}/h}u_1|^2_{L^2(S)} + h^3|\mathrm{e}^{\varphi_{r,1}/h}\nabla u_1|^2_{L^2(S)} + h|\mathrm{e}^{\varphi_{r,1}/h}\phi_1|^2_{L^2(\Gamma_1)} \\ +h^3|\mathrm{e}^{\varphi_{r,1}/h}\phi_2|^2_{L^2(\Gamma_1)} + h^{-1}|\mathrm{e}^{\varphi_{r,1}/h}a\partial_\nu u_1|^2_{L^2(\Gamma_1)} + h|\mathrm{e}^{\varphi_{r,1}/h}bu_1|^2_{L^2(\Gamma_1)}). \end{aligned} \tag{5.12}$$

We addition the two last estimates for $r, k \in \{1, 2\}$ and using the properties of phases ($\varphi_{r,k} < \varphi_{\sigma(r),k}$ in $\bigcup\limits_{j_r=1}^{N_{r,k}} B(x^k_{1,j_r}, 2\epsilon)$) then we can absorb the terms $[\Delta, \chi_{r,1}]y_1$ and $[\Delta, \chi_{r,2}]y_2$ at the right hand side of (5.12) into the le left hand side for $h > 0$ small. Morever by the expression of y_1 and y_2 in (5.6) we obtain

$$\begin{aligned} h\int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\Delta u_1|^2\mathrm{d}x + h\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h} + \mathrm{e}^{2\varphi_{2,2}/h}\right)|u_2|^2\mathrm{d}x \\ +h^3\int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\nabla y_1|^2\mathrm{d}x + h^3\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h} + \mathrm{e}^{2\varphi_{2,2}/h}\right)|\nabla u_2|^2\mathrm{d}x \\ +h^3\int_S\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\partial_\nu u_2|^2\mathrm{d}x \leq C\Bigg(h^4\int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\Phi_1|^2\mathrm{d}x \\ +h^4\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h} + \mathrm{e}^{2\varphi_{2,2}/h}\right)|\Phi_2|^2\mathrm{d}x + \int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|u_1|^2\mathrm{d}x \\ +h\int_S\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|u_1|^2\mathrm{d}x + h^3\int_S\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\nabla u_1|^2\mathrm{d}x \\ +h\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\phi_1|^2\mathrm{d}x + h^{-1}\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)|\phi_2|^2\mathrm{d}x \\ +h^{-1}\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)a|\partial_\nu u_1|^2\mathrm{d}x + h\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h} + \mathrm{e}^{2\varphi_{2,1}/h}\right)b|u_1|^2\mathrm{d}x\Bigg). \end{aligned} \tag{5.13}$$

We will now focus to the system (5.8). Let $\psi_{1,1}$, $\psi_{2,1}$, $\psi_{1,2}$ and $\psi_{2,2}$ four phases as described in section 4. Let $\chi_{1,1}$, $\chi_{2,1}$, $\chi_{1,2}$ and $\chi_{2,2}$ four cutt-off functions equal to one respectively in $\left(\bigcup\limits_{l_1=1}^{M_{1,1}} B(y^1_{1,l_1}, 2\epsilon)\right)^c$, $\left(\bigcup\limits_{l_2=1}^{M_{2,1}} B(y^1_{2,l_2}, 2\epsilon)\right)^c$, $\left(\bigcup\limits_{l_1=1}^{M_{1,2}} B(y^2_{1,l_1}, 2\epsilon)\right)^c$ and $\left(\bigcup\limits_{l_2=1}^{M_{2,2}} B(y^2_{2,l_2}, 2\epsilon)\right)^c$ and supported in $\left(\bigcup\limits_{l_1=1}^{M_{1,1}} B(y^1_{1,l_1}, \epsilon)\right)^c$, $\left(\bigcup\limits_{l_2=1}^{M_{2,1}} B(y^1_{2,l_2}, \epsilon)\right)^c$, $\left(\bigcup\limits_{l_1=1}^{M_{1,2}} B(y^2_{1,l_1}, \epsilon)\right)^c$ and $\left(\bigcup\limits_{l_2=1}^{M_{2,2}} B(y^2_{2,l_2}, \epsilon)\right)^c$ respectively (in order to eliminate the critical points of the phases functions $\psi_{1,1}$, $\psi_{2,1}$, $\psi_{1,2}$ and $\psi_{2,2}$).

Then for $r = 1, 2$, where we are noting by $\tilde{z}_{r,1} = \chi_{r,1}z_1$ and $\tilde{z}_{r,2} = \chi_{r,1}z_2$, and we obtain from the system (5.8) the following equation

$$\left\{\begin{array}{lll} \Delta\tilde{z}_{r,1}+\dfrac{1}{h}\tilde{z}_{r,1}=\widetilde{\Psi}_{r,1} & \text{in} & \Omega_1 \\ \Delta\tilde{z}_{r,2}-\dfrac{1}{h}\tilde{z}_{r,2}=\widetilde{\Psi}_{r,2} & \text{in} & \Omega_1 \\ h\tilde{z}_{r,1}=\tilde{z}_{r,2} & \text{on} & S \\ h\partial_\nu\tilde{z}_{r,1}+\partial_\nu\tilde{z}_{r,2}=h\partial_\nu u_2 & \text{on} & S \\ \tilde{z}_{r,1}=\phi_1-i\mu a\partial_\nu u_1-|\mu|u_1 & \text{on} & \Gamma_1 \\ \partial_\nu\tilde{z}_{r,1}=\phi_2+i\mu b u_1-|\mu|\partial_\nu u_1 & \text{on} & \Gamma_1, \end{array}\right. \tag{5.14}$$

where

$$\left\{\begin{array}{ll} \widetilde{\Psi}_{r,1}=[\Delta,\chi_{r,1}]z_1+\chi_{r,1}.\Phi_1 & \text{in } \Omega_1 \\ \widetilde{\Psi}_{r,2}=[\Delta,\chi_{r,2}]z_2+\chi_{r,2}.z_1 & \text{in } \Omega_1. \end{array}\right. \tag{5.15}$$

Applying now Carleman estimate of Theorem 4.2 to the system (5.14) then for $r=1,2$ and by assumption (1.2) we obtain

$$\begin{aligned} h^3\|\mathrm{e}^{\psi_{r,1}/h}\tilde{z}_{r,1}\|^2_{L^2(V_{r,1})}+h\|\mathrm{e}^{\psi_{r,2}/h}\tilde{z}_{r,2}\|^2_{L^2(V_{r,2})}+h^5\|\mathrm{e}^{\psi_{r,1}/h}\nabla\tilde{z}_{r,1}\|^2_{L^2(V_{r,1})}+ \\ h^3\|\mathrm{e}^{\psi_{r,2}/h}\nabla\tilde{z}_{r,2}\|^2_{L^2(V_{r,2})}+h|\mathrm{e}^{\psi_{r,2}/h}\tilde{z}_{r,2}|^2_{L^2(S)}+h^3|\mathrm{e}^{\psi_{r,2}/h}\nabla\tilde{z}_{r,2}|^2_{L^2(S)} \\ \le C(h^6\|\mathrm{e}^{\psi_{r,1}/h}\widetilde{\Psi}_{r,1}\|^2_{L^2(V_{r,1})}+h^4\|\mathrm{e}^{\psi_{r,2}/h}\widetilde{\Psi}_{r,2}\|^2_{L^2(V_{r,2})}+h^5|\mathrm{e}^{\psi_{r,1}/h}\partial_\nu u_2|^2_{L^2(S)}+ \\ h|\mathrm{e}^{\psi_{r,1}/h}\phi_1|^2_{L^2(\Gamma_1)}+h^3|\mathrm{e}^{\psi_{r,1}/h}\phi_2|^2_{L^2(\Gamma_1)}+h^{-1}|\mathrm{e}^{\psi_{r,1}/h}a\partial_\nu u_1|^2_{L^2(\Gamma_1)}+h^{-1}|\mathrm{e}^{\psi_{r,1}/h}bu_1|^2_{L^2(\Gamma_1)}). \end{aligned}$$

Which by (5.15) we find

$$\begin{aligned} h^3\|\mathrm{e}^{\psi_{r,1}/h}\tilde{z}_{r,1}\|^2_{L^2(V_{r,1})}+h\|\mathrm{e}^{\psi_{r,2}/h}\tilde{z}_{r,2}\|^2_{L^2(V_{r,2})}+h^5\|\mathrm{e}^{\psi_{r,1}/h}\nabla\tilde{z}_{r,1}\|^2_{L^2(V_{r,1})}+ \\ h^3\|\mathrm{e}^{\psi_{r,2}/h}\nabla\tilde{z}_{r,2}\|^2_{L^2(V_{r,2})}+h|\mathrm{e}^{\psi_{r,2}/h}u_1|^2_{L^2(S)}+h^3|\mathrm{e}^{\psi_{r,2}/h}\nabla u_1|^2_{L^2(S)} \\ \le C(h^6\|\mathrm{e}^{\psi_{r,1}/h}\Phi_1\|^2_{L^2(V_{r,1})}+h^4\|\mathrm{e}^{\psi_{r,2}/h}z_1\|^2_{L^2(V_{r,2})}+h^5|\mathrm{e}^{\psi_{r,1}/h}\partial_\nu u_2|^2_{L^2(S)}+ \\ h^6\|\mathrm{e}^{\psi_{r,1}/h}[\Delta,\chi_{r,1}]z_1\|_{L^2(U_{r,1})}+h^4\|\mathrm{e}^{\psi_{r,2}/h}[\Delta,\chi_{r,2}]z_2\|_{L^2(U_{r,2})}+h|\mathrm{e}^{\psi_{r,1}/h}\phi_1|^2_{L^2(\Gamma_1)}+ \\ h^3|\mathrm{e}^{\psi_{r,1}/h}\phi_2|^2_{L^2(\Gamma_1)}+h^{-1}|\mathrm{e}^{\psi_{r,1}/h}a\partial_\nu u_1|^2_{L^2(\Gamma_1)}+h^{-1}|\mathrm{e}^{\psi_{r,1}/h}bu_1|^2_{L^2(\Gamma_1)}). \end{aligned} \tag{5.16}$$

For the same reasons as previously we can absorb the terms $[\Delta,\chi_{r,1}]z_1$ and $[\Delta,\chi_{r,2}]z_2$ at the right hand side of (5.16) to the left hand side by adding the two last estimates, namely we get

$$\begin{aligned} h^3\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|z_1|^2\mathrm{d}x+h\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|z_2|^2\mathrm{d}x \\ +h^5\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\nabla z_1|^2\mathrm{d}x+h^3\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\nabla z_2|^2\mathrm{d}x \\ +h\int_S\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|u_1|^2\mathrm{d}x+h^3\int_S\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\nabla u_1|^2\mathrm{d}x \\ \le C\Bigg(h^6\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\Phi_1|^2\mathrm{d}x+h^4\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|z_1|^2\mathrm{d}x \\ +h\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_1|^2\mathrm{d}x+h^3\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_2|^2\mathrm{d}x \\ +h^{-1}\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)a|\partial_\nu u_1|^2\mathrm{d}x+h^{-1}\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)b|u_1|^2\mathrm{d}x \\ +h^5\int_S\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\partial_\nu u_2|^2\mathrm{d}x\Bigg). \end{aligned} \tag{5.17}$$

Then by the expression of z_1 and z_2 in (5.8) we can wite (5.17) after multiplying by h^{-1} in the following form

$$\begin{aligned}
&h^2\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|z_1|^2\mathrm{d}x+\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|u_1|^2\mathrm{d}x\\
&+h^4\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\nabla z_1|^2\mathrm{d}x+h^2\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\nabla z_2|^2\mathrm{d}x\\
&+\int_{S}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|u_1|^2\mathrm{d}x+h^2\int_{S}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\nabla u_1|^2\mathrm{d}x\\
&\le C\Bigg(h^5\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\Phi_1|^2\mathrm{d}x+h^3\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\Delta u_1|^2\mathrm{d}x\\
&+\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_1|^2\mathrm{d}x+h^2\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_2|^2\mathrm{d}x\\
&+h^{-2}\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)a|\partial_\nu u_1|^2\mathrm{d}x+h^{-2}\int_{\Gamma_1}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)b|u_1|^2\mathrm{d}x\\
&+h^4\int_{S}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\partial_\nu u_2|^2\mathrm{d}x\Bigg).
\end{aligned}\tag{5.18}$$

By making the choise of the weight function as described in section 4 (in one hand $\varphi_{1,1}=\psi_{1,2}$ and $\varphi_{2,1}=\psi_{2,2}$ in Ω_1 and in an other hand $\varphi_{1,1}=\psi_{1,1}$ and $\varphi_{2,1}=\psi_{2,1}$ on the edge S) we get by combining (5.13) and (5.18) that

$$\begin{aligned}
&h\int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}\right)|\Delta u_1|^2\mathrm{d}x+h\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h}+\mathrm{e}^{2\varphi_{2,2}/h}\right)|u_2|^2\mathrm{d}x\\
&+h^3\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h}+\mathrm{e}^{2\varphi_{2,2}/h}\right)|\nabla u_2|^2\mathrm{d}x+h^3\int_{S}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}\right)|\partial_\nu u_2|^2\mathrm{d}x\\
&\le C\Bigg(h^4\int_{\Omega_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}+\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\Phi_1|^2\mathrm{d}x\\
&+h^4\int_{\Omega_2}\left(\mathrm{e}^{2\varphi_{1,2}/h}+\mathrm{e}^{2\varphi_{2,2}/h}\right)|\Phi_2|^2\mathrm{d}x\\
&+\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}+\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_1|^2\mathrm{d}x\\
&+h^{-1}\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}+\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\phi_2|^2\mathrm{d}x\\
&+\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}+\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)a|\partial_\nu u_1|^2\mathrm{d}x\\
&+\int_{\Gamma_1}\left(\mathrm{e}^{2\varphi_{1,1}/h}+\mathrm{e}^{2\varphi_{2,1}/h}+\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)b|u_1|^2\mathrm{d}x\\
&+h^3\int_{\Omega_1}\left(\mathrm{e}^{2\psi_{1,2}/h}+\mathrm{e}^{2\psi_{2,2}/h}\right)|\Delta u_1|^2\mathrm{d}x+h^4\int_{S}\left(\mathrm{e}^{2\psi_{1,1}/h}+\mathrm{e}^{2\psi_{2,1}/h}\right)|\partial_\nu u_2|^2\mathrm{d}x\Bigg).
\end{aligned}\tag{5.19}$$

Then we can absorb the two last terms in the left hand side of (5.19) to the left hand side for h small to get

$$\begin{aligned}
\|\Delta u_1\|^2_{L_2(\Omega_1)}+\|u_2\|^2_{L_2(\Omega_2)}+\|\nabla u_2\|^2_{L_2(\Omega_2)}&\le C\mathrm{e}^{C/h}\Big(\|\Phi_1\|^2_{L^2(\Omega_1)}+\|\Phi_2\|^2_{L^2(\Omega_2)}+\\
&|\phi_1|^2_{L^2(\Gamma_1)}+|\phi_2|^2_{L^2(\Gamma_1)}+\int_{\Gamma_1}a|\partial_\nu u_1|^2\mathrm{d}x+\int_{\Gamma_1}b|u_1|^2\mathrm{d}x\Big).
\end{aligned}\tag{5.20}$$

So that the combination of (5.5) and (5.20) give us

$$\begin{aligned}\|\Delta u_1\|^2_{L^2(\Omega_1)} + \|u_2\|^2_{L_2(\Omega_2)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} \leq C\mathrm{e}^{C/h}\Big(\|\Phi_1\|^2_{L^2(\Omega_1)} + \|\Phi_2\|^2_{L^2(\Omega_2)} +\\ \|\Phi_1\|_{L^2(\Omega_1)}.\|u_1\|_{L^2(\Omega_1)} + \|\Phi_2\|_{L^2(\Omega_2)}.\|u_2\|_{L^2(\Omega_2)} + |\phi_1|^2_{L^2(\Gamma_1)} + |\phi_2|^2_{L^2(\Gamma_1)}\\ +|\phi_1|_{L^2(\Gamma_1)}.|\partial_\nu u_1|_{L^2(\Gamma_1)} + |\phi_2|_{L^2(\Gamma_1)}.|u_1|_{L^2(\Gamma_1)}\Big).\end{aligned}$$

By Young inequality, Remark 2.1 and expressions of ϕ_1 and ϕ_2 in (5.2) we get

$$\|\Delta u_1\|^2_{L^2(\Omega_1)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} \leq C\mathrm{e}^{C/h}\Big(\|\Phi_1\|^2_{L^2(\Omega_1)} + \|\Phi_2\|^2_{L^2(\Omega_2)} + \|f_1\|^2_{H^2(\Omega_1)}\Big).$$

Using the expressions of Φ_1 and Φ_2 in (5.2) we find

$$\|\Delta u_1\|^2_{L^2(\Omega_1)} + \|\nabla u_2\|^2_{L^2(\Omega_2)} \leq C\mathrm{e}^{C/h}\Big(\|F\|^2_X + \|G\|^2_H\Big). \tag{5.21}$$

By (5.21) and the expression of v in (5.3) we get

$$\|v\|^2_H \leq \|F\|^2_X + |\mu|^2.\|u\|^2_{L^2(\Omega)} \leq C\mathrm{e}^{C/h}\Big(\|F\|^2_X + \|G\|^2_H\Big).$$

Which by (5.21) give us

$$\|(u,v)\|^2_{\mathcal{H}} \leq C\mathrm{e}^{C/h}\Big(\|F\|^2_X + \|G\|^2_H\Big).$$

In other words we have

$$\|(u,v)\|^2_{\mathcal{H}} \leq C\mathrm{e}^{C/h}\|(\mathcal{A} - i\mu\,\mathrm{Id})(u,v)\|_{\mathcal{H}}.$$

Hence $(\mathcal{A} - i\mu\mathrm{Id})$ is injective then bijective in $\mathcal{D}(\mathcal{A})$ and we have

$$\|(\mathcal{A} - i\mu\,\mathrm{Id})^{-1}\|_{\mathcal{L}(\mathcal{H})} \leq C\mathrm{e}^{C|\mu|},$$

for $|\mu|$ large enough.

A Appendix: Semi Classical analysis

The aim of this section is to give some results from the semi-classical analysis which is an indispensable means to prove the local Carleman estimates. We shall use of the notation $\langle\xi\rangle = (1+|\xi|^2)^{\frac{1}{2}}$. We denote by S^m the space of smooth functions $a(x,\xi,h)$ defined for h a semi-classical parameter in $]0,h_0]$, that satisfy the following property:

$$\forall\,\alpha,\beta \in \mathbb{N}^n,\ \exists\, C_{\alpha,\beta} > 0 \quad |\partial_x^\alpha \partial_\xi^\beta a(x,\xi,h)| \leq C_{\alpha,\beta}\langle\xi\rangle^{m-\beta}; \quad \forall\, x,\xi \in \mathbb{R}^n,\ \forall\, h \in]0,h_0].$$

We define Ψ^m as the space of pseudo-differential operators (PDO) A; for $a \in S^m$ and $u \in S(\mathbb{R}^n)$ by

$$Au(x) := \mathrm{Op}(a)(u)(x) = \frac{1}{(2\pi h)^n}\iint \mathrm{e}^{\frac{i}{h}\langle x-y,\xi\rangle} a(x,\xi,h)u(y)\,\mathrm{d}y\,\mathrm{d}\xi.$$

For A a PDO we denote by $\sigma(A)$ his principal symbol. Our analysis is based on the use of the pseudo-diffential calculs thechniques, such as constraction of parametrices, composition formula, adjoint formula, etc. (we refert the reader to [Hör85, Mar02]), in particular for $a \in S^m$ and $b \in S^{m'}$ we have

- $\mathrm{Op}(a) \circ \mathrm{Op}(b) \in \Psi^{m+m'}$ such that

$$(a\sharp b)(x,\xi,h) \sim \sum_{|\alpha|\geq 0} \frac{h^{|\alpha|}}{i^{|\alpha|}\alpha!}\partial_\xi^\alpha a(x,\xi,h)\partial_x^\alpha b(x,\xi,h), \quad \alpha! = \alpha_1! \dots \alpha_n!$$

- In particular, $[\mathrm{Op}(a), \mathrm{Op}(b)] \in h\Psi^{m+m'-1}$ and we have

$$\sigma([\mathrm{Op}(a), \mathrm{Op}(b)]) = \frac{h}{i}\{a, b\} = \frac{h}{i}\sum_{j=1}^{n} \partial_{\xi_j} a(x,\xi,h)\partial_{x_j} b(x,\xi,h) - \partial_{x_j} a(x,\xi,h)\partial_{\xi_j} b(x,\xi,h).$$

- $\mathrm{Op}(a)^* = \mathrm{Op}(c) \in \Psi^m$ such that $\sigma(\mathrm{Op}(a)^*) = \overline{a}$ and

$$c(x,\xi,h) \sim \sum_{|\alpha|\geq 0} \frac{h^{|\alpha|}}{i^{\alpha}\alpha!}\partial_x^{\alpha}\partial_{\xi}^{\alpha}\overline{a(x,\xi,h)}.$$

Now we introduce tangential symbols and associated operators. We set $x = (x', x_n)$; $x' = (x_1, \dots, x_{n-1})$ and we denote by S_t^m the space of smooth functions $a(x,\xi',h)$ defined for a small parameter $h \in]0, h_0]$ that satisfy the following property:
$\forall\, \alpha \in \mathbb{N}^n, \beta \in \mathbb{N}^{n-1}$, $\exists\, C_{\alpha,\beta} > 0$ such that

$$|\partial_x^{\alpha}\partial_{\xi'}^{\beta} a(x,\xi',h)| \leq C_{\alpha,\beta}\langle\xi'\rangle^{m-\beta}; \quad \forall\, x \in \mathbb{R}^n,\ \xi' \in \mathbb{R}^{n-1},\ \forall\, h \in]0, h_0].$$

As above we define Ψ_t^m as the space of tangential pseudo-differential operators (TPDO) A; for $a \in S_t^m$ and $u \in S(\mathbb{R}^n)$ by

$$Au(x) := \mathrm{op}_t(a)(u)(x) = \frac{1}{(2\pi h)^{n-1}}\iint \mathrm{e}^{\frac{i}{h}\langle x'-y',\xi'\rangle} a(x,\xi',h)u(y',x_n)\,\mathrm{d}y'\,\mathrm{d}\xi'.$$

For $a \in S_t^m$, $b \in S_t^{m'}$ the composition formula for the tangential symbols is given by

$$(a\sharp b)(x,\xi',h) \sim \sum_{|\alpha|\geq 0} \frac{h^{|\alpha|}}{i^{|\alpha|}\alpha!}\partial_{\xi'}^{\alpha} a(x,\xi',h)\partial_x^{\alpha} b(x,\xi',h) \in S_t^{m+m'}.$$

We denote by $\mathcal{D}_t^m$ the sub-space of Ψ_t^m, where $m \in \mathbb{N}$, constitute of operators whose symbols are polynomial relative to the variable ξ'. We denote by Λ^s (resp. Λ_t^s) the PDO (resp. TPDO) whose symbole is $\langle\xi\rangle^s$ (resp. $\langle\xi'\rangle^s$) for $s \in \mathbb{R}$ and we introduce the norme $\|u\|_s = \|\Lambda^s u\|_0$ and the Sobolev space

$$\mathcal{H}^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) :\ \|u\|_s < \infty\}.$$

We shall denote by $(\,.\,,\,.\,)_0$ the inner product for functions defined in $\{x_n = 0\}$ by $(f,g)_0 = \langle f,g\rangle_{L^2(\mathbb{R}^{n-1})}$ and the induced norme is denoted by $|f|_0$. For $s \in \mathbb{R}$, we introduce $|u|_s = |\Lambda_t^s u|_0$. Follow to [Mar02] we have the following resut of continuity of the PDO.

Proposition A.1 *For $s \in \mathbb{R}$; if $a \in S^m$, then $\mathrm{Op}(a) : \mathcal{H}^s \longrightarrow \mathcal{H}^{s-m}$ continuously, uniformly in h.*

Follow to [RR10, Lemma 2.10] we have this technical result

Lemma A.1 *If $a \in S_t^m$ and $A \in S^{-\infty}$ such that $\mathrm{supp}(a) \cap \mathrm{supp}(A) = \emptyset$. Then we have*

$$\mathrm{op}_t(a) \circ \mathrm{Op}(A) \in \bigcap_{N\in\mathbb{N}} h^N\Psi^{-N} \quad \textit{and} \quad \mathrm{Op}(A) \circ \mathrm{op}_t(a) \in \bigcap_{N\in\mathbb{N}} h^N\Psi^{-N}.$$

Remark A.1 According to [LR97, p.486] we have for all $j \in \mathbb{N}$, there existe $C > 0$ such that

$$|u_{|x_n=0}|_j \leq Ch^{-\frac{1}{2}}\|u\|_{j+1} \quad \forall\, u \in \mathscr{C}_c^{\infty}(\mathbb{R}^n).$$

We finish this part by the Gårding lemma which is an importing tool for the proof of Carleman estimates.

Lemma A.2 *If $A \in \Psi^m$ verifying that for some $C > 0$*

$$\mathrm{Re}(\sigma(A))(x,\xi,h) \geq C\langle\xi\rangle^m \quad \forall\, x \in K, \forall\, \xi \in \mathbb{R}^n, \forall\, h \in]0, h_0]$$

where K is a compact subset in $\mathbb{R}^n$. Then there existe $C' > 0$ such that for $h_1 > 0$ small enough we have

$$\mathrm{Re}\langle \mathrm{Op}(a)u, u\rangle_{L^2(\mathbb{R}^n)} \geq C'\|u\|_{\frac{m}{2}}^2 \quad \forall\, u \in \mathscr{C}_c^{\infty}(K),\ \forall\, h \in]0, h_1].$$

References

[AN10] K. Ammari and S. Nicaise. Stabilization of a transmission wave/plate equation. *Journal of Differential Equations*, 249:707–727, 2010.

[AV09] K. Ammari and G. Vodev. Boundary stabilization of the transmission problem for the Bernoulli-Euler plate equation. *CUBO a mathematical journal*, 11:39–49, 2009.

[BD08] C.J.K Batty and T. Duyckaerts. Non-uniform stability for bounded semi-groups on Banach spaces. *Journal of Evolution Equation*, pages 765–780, 2008.

[Bel03] M. Bellassoued. Carleman estimates and distribution of resonnances for the transparent obstace and application to the stabilization. *Asymptotic Anal.*, 35:257–279, 2003.

[Bur98] N. Burq. Décroissance de l'énergie locale de l'équation des ondes pour le problème extérieur et absence de résonnance au voisinage du réel. *Acta Math.*, 180:1–29, 1998.

[Duy07] T. Duyckaerts. Optimal decay rates of the energy of a hyperbolic-parabolic system coupled by an interface. *Asymptot. Anal.*, 51:17–45, 2007.

[EMT04] Y. Eidelman, V. Milman, and A. Tsolomitis. *Functional Analysis An Introduction*. American Mathematical Society, 2004.

[Fat11] I.K. Fathallah. Logarithmic decay of the energy for an hyperbolic-parabolic coupled system. *ESAIM-control Optimisation and Calculus of Variations*, 17:801–835, 2011.

[Has13] F. Hassine. Weak stabilization of a transmission Bernoulli-Euler plate equation with force and moment feedback. *arXiv:1301.3536[math.AP]*, 2013.

[Hör63] L. Hörmander. *Linear partial diffenrential operators*. Springer-verlag, 1963.

[Hör85] L. Hörmander. *The analysis of linear partial diffenrential operators*, volume 3. Springer-verlag, 1985.

[LR95] G. Lebeau and L. Robbiano. Contrôle exacte de l'équation de la chaleur. *Comm. Partial Differential Equations*, 20:335–356, 1995.

[LR97] G. Lebeau and L. Robbiano. Stabilisation de l'équation des ondes par le bord. *Duke mathematical journal*, 86:465–491, december 1997.

[Mar02] A. Martinez. *An introduction to semiclassical and microlocal analysis*. Springer-verlag, 2002.

[RR10] J. Le Rousseau and L. Robbiano. Carleman estimate for elliptic operators with coefficients with jumps at an interface in arbitrary dimension and application to the null controllability of linear parabolic equations. *Arch. Rational Mech. Anal.*, 195:953–990, 2010.

[RZZ05] J. Rauch, X. Zhang, and E. Zuazua. Polynomial decay for hyperbolic-parabolic coupled system. *Math. Pures Appl.*, 84:407–470, 2005.

[Teb12] L. Tebou. Energy decay estimates for some weakly coupled Euler-Bernoulli and wave equations with indirect damping mechanisms. *Mathematical control and related fields*, 2(1):45–60, 2012.

[TW09] M. Tucsnak and G. Weiss. *Observation and control for operator semigroups*. Birkhäuser Verlag AG, 2009.

[ZZ06] X. Zhang and E. Zuazua. Asyptotic behavior of a hyperbolic-parabolic coupled system arising in fluid-structure interaction. *International Series of Numerical Mathematics*, 154:445–455, 2006.

[ZZ07] X. Zhang and E. Zuazua. Long-time behavior of a coupled heat-wave system arising in fluid-structure interaction. *Arch. Ration. Mech. Anal*, 184:49–120, 2007.